

RC
268.5
1155
no. 120
1979

National Cancer Institute
CARCINOGENESIS
Technical Report Series
NO. 120
1979

**BIOASSAY OF
PIPERONYL BUTOXIDE
FOR POSSIBLE CARCINOGENICITY**

CAS No. 51-03-6

NCI-CG-TR-120

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health



Library
National Archives at Western
College, Maryland 20014

BIOASSAY OF
PIPERONYL BUTOXIDE
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
Division of Cancer Cause and Prevention
U.S. National Cancer Institute
National Institutes of Health
Bethesda, Maryland 20014

Technical report series

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health

RC
268.5
455
no. 120
1979

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

National Institutes of Health

REPORT ON BIOASSAY OF PARATHION FOR POSSIBLE CARCINOGENICITY

Availability

Parathion (CAS 56-38-2) has been tested for cancer-causing activity with rats and mice in the Bioassay Program, Division of Cancer Cause and Prevention, National Cancer Institute. A report is available to the public.

Summary: A bioassay for possible carcinogenicity of technical-grade parathion was conducted by administering the test chemical in the diet to Osborne-Mendel rats and B6C3F1 mice. Applications of the chemical include use as a pesticide.

It is concluded that under the conditions of this bioassay, parathion was not carcinogenic to B6C3F1 mice. In the male and female Osborne-Mendel rats receiving parathion in their diet, there was a higher incidence of cortical tumors of the adrenal than in pooled or historical controls, suggesting that parathion is carcinogenic to this strain of rat.

Single copies of the report are available from the Office of Cancer Communications, National Cancer Institute, Building 31, Room 10A21, National Institutes of Health, Bethesda, Maryland 20014.

Dated:
November 16, 1978

Director
National Institutes of Health

(Catalogue of Federal Domestic Assistance Program Number 13.393, Cancer Cause and Prevention Research)



BIOASSAY OF
PIPERONYL BUTOXIDE
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
Division of Cancer Cause and Prevention
National Cancer Institute
National Institutes of Health

FOREWORD: This report presents the results of the bioassay of piperonyl butoxide conducted for the Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute (NCI), National Institutes of Health, Bethesda, Maryland. This is one of a series of experiments designed to determine whether selected chemicals have the capacity to produce cancer in animals. Negative results, in which the test animals do not have a greater incidence of cancer than control animals, do not necessarily mean that the test chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of circumstances. Positive results demonstrate that the test chemical is carcinogenic for animals under the conditions of the test and indicate that exposure to the chemical is a potential risk to man. The actual determination of the risk to man from animal carcinogens requires a wider analysis.

CONTRIBUTORS: This bioassay of piperonyl butoxide was conducted by Frederick Cancer Research Center (FCRC) (1), Frederick, Maryland, operated for NCI (2) by Litton Bionetics, Inc., Kensington, Maryland (3).

The manager of the bioassay at FCRC was Dr. D. Creasia. The program manager was Dr. B. Ulland, and the toxicologist was Dr. E. Gordon. Ms. S. Toms was responsible for management of data, Mr. D. Cameron for management of histopathology, Mr. L. Callahan for management of the computer branch, and Mr. R. Cypher for the management of the facilities. Mr. A. Butler performed the computer services. Histopathologic evaluations for rats were performed by Dr. R. A. Renne (4,5), and the histopathologic evaluations for mice were performed by Dr. C. E. Gilmore (4). The diagnoses included in this report represent the interpretations of Drs. Renne and Gilmore.

Animal pathology tables and survival tables were compiled at EG&G Mason Research Institute (6). Statistical analyses were performed by Dr. J. R. Joiner (7) and Ms. P. L. Yong (7), using methods selected for the bioassay program by Dr. J. J. Gart (8).

The chemicals used in this bioassay were analyzed at Frederick Cancer Research Center by Dr. W. Zielinsky (1). The results of these analyses were reviewed by Dr. C. W. Jameson (7) and Ms. P. M. Wagner (7).

This report was prepared at Tracor Jitco (7) under the direction of NCI. Those responsible for the report at Tracor Jitco were Dr. L. A. Campbell, Director of the Bioassay Program; Dr. S. S. Olin, Deputy Director for Science; Dr. J. F. Robens, toxicologist; Dr. R. L. Schueler, pathologist; Dr. G. L. Miller, Ms. M. S. King, Ms. L. A. Waitz, and Mr. W. D. Reichardt, bioscience writers; and Dr. E. W. Gunberg, technical editor, assisted by Ms. Y. E. Presley and Ms. P. J. Graboske.

The following scientists at NCI (2) were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. Kenneth C. Chu, Dr. Cipriano Cueto, Jr., Dr. J. Fielding Douglas, Dr. Dawn G. Goodman (9), Dr. Richard A. Griesemer, Dr. Morton H. Levitt, Dr. Harry A. Milman, Dr. Thomas W. Orme, Dr. Robert A. Squire (10), Dr. Sherman F. Stinson, Dr. Jerrold M. Ward, and Dr. Carrie E. Whitmire.

-
- (1) Frederick Cancer Research Center, P.O. Box B, Frederick, Maryland.
 - (2) Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
 - (3) Litton Bionetics, Inc., 5516 Nicholson Lane, Kensington, Maryland.
 - (4) Experimental Pathology Laboratories, Inc., P.O. Box 474, Herndon, Virginia.
 - (5) Now with Battelle Pacific Northwest Laboratories, Battelle Boulevard, Richland, Washington.

- (6) EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.
- (7) Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.
- (8) Mathematical Statistics and Applied Mathematics Section, Biometry Branch, Field Studies and Statistics, Division of Cancer Cause and Prevention, National Cancer Institute National Institutes of Health, Bethesda, Maryland.
- (9) Now with Clement Associates, Inc., 1010 Wisconsin Avenue, N.W., Suite 660, Washington, D. C.
- (10) Now with the Division of Comparative Medicine, Johns Hopkins University, School of Medicine, Traylor Building, Baltimore, Maryland.

1. The first part of the paper discusses the importance of the study and the objectives of the research. It also provides a brief overview of the literature review and the methodology used in the study.

2. The second part of the paper presents the results of the study. It includes a detailed analysis of the data collected and the findings of the research. The results are presented in a clear and concise manner, with appropriate use of tables and figures.

3. The third part of the paper discusses the implications of the study and the conclusions drawn from the research. It also provides a brief summary of the key findings and the overall contribution of the study to the field.

4. The fourth part of the paper provides a detailed discussion of the limitations of the study and the areas for future research. It also includes a brief discussion of the ethical considerations and the potential impact of the study on society.

5. The fifth part of the paper provides a detailed discussion of the conclusions drawn from the research. It also includes a brief summary of the key findings and the overall contribution of the study to the field.

6. The sixth part of the paper provides a detailed discussion of the implications of the study and the conclusions drawn from the research. It also provides a brief summary of the key findings and the overall contribution of the study to the field.

7. The seventh part of the paper provides a detailed discussion of the limitations of the study and the areas for future research. It also includes a brief discussion of the ethical considerations and the potential impact of the study on society.

8. The eighth part of the paper provides a detailed discussion of the conclusions drawn from the research. It also includes a brief summary of the key findings and the overall contribution of the study to the field.

9. The ninth part of the paper provides a detailed discussion of the implications of the study and the conclusions drawn from the research. It also provides a brief summary of the key findings and the overall contribution of the study to the field.

10. The tenth part of the paper provides a detailed discussion of the limitations of the study and the areas for future research. It also includes a brief discussion of the ethical considerations and the potential impact of the study on society.

SUMMARY

A bioassay of technical-grade piperonyl butoxide for possible carcinogenicity was conducted by administering the test chemical in feed to Fischer 344 rats and B6C3F1 mice.

Groups of 50 rats of each sex were administered piperonyl butoxide in the diet at one of two doses, either 5,000 or 10,000 ppm, for 107 weeks. Matched controls consisted of 20 untreated rats of each sex. All surviving rats were killed at the end of the period of administration of the test chemical.

Groups of 50 mice of each sex were initially administered piperonyl butoxide at one of two doses, either 2,500 or 5,000 ppm. After week 30, the doses for the mice were reduced to 500 and 2,000 ppm, respectively, and administration of the test chemical at the lowered doses was continued for 82 weeks. The time-weighted average doses for the mice were either 1,036 or 2,804 ppm. Matched controls consisted of 20 untreated mice of each sex. All surviving mice were killed at the end of the period of administration of the test chemical.

Mean body weights of dosed groups of rats and mice of each sex were lower than those of corresponding control groups, and the depressions in body weights were dose related. Survival of the rats and mice was unaffected by the piperonyl butoxide and was 80% or greater in all groups at week 90 of the bioassay; thus, sufficient numbers of dosed and control rats and mice of each sex were at risk for the development of late-appearing tumors.

In the female rats, lymphomas occurred at incidences that were dose related ($P = 0.007$); in a direct comparison, the incidence of the tumor in the high-dose group was higher ($P = 0.020$) than that in the control group (controls 1/20, low-dose 7/50, high-dose 15/50). However, the incidence of lymphomas, leukemias, and reticuloses in historical-control female Fischer 344 rats at the same laboratory was 19/191 (10%). These historical-control groups include one with an incidence of animals with lymphoma or leukemia of 7/20 (35%) and another with an incidence of 6/20 (30%). Thus, the incidence of lymphomas in the control female rats of the present bioassay may have been abnormally low, and the occurrence of the higher incidence in the dosed groups cannot be clearly related to administration of piperonyl butoxide.

In the male mice, adenomas of the lacrimal gland occurred at incidences that were dose related ($P = 0.023$), but in direct comparisons the incidences in the individual dosed groups were not significantly higher than that in the control group (controls 0/20, low-dose 0/49, high-dose 4/50); thus, the occurrence of this tumor in the male mice was not clearly related to administration of the test chemical.

It is concluded that under the conditions of this bioassay, piperonyl butoxide was not carcinogenic for Fischer 344 rats or B6C3F1 mice.

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction	1
II. Materials and Methods	3
A. Chemical	3
B. Dietary Preparation	4
C. Animals	4
D. Animal Maintenance	5
E. Subchronic Studies	7
F. Chronic Studies	10
G. Clinical and Pathologic Examinations	13
H. Data Recording and Statistical Analyses	14
III. Results - Rats	21
A. Body Weights and Clinical Signs (Rats)	21
B. Survival (Rats)	21
C. Pathology (Rats)	24
D. Statistical Analyses of Results (Rats)	31
IV. Results - Mice	33
A. Body Weights and Clinical Signs (Mice)	33
B. Survival (Mice)	33
C. Pathology (Mice)	36
D. Statistical Analyses of Results (Mice)	38
V. Discussion	41
VI. Bibliography	45

APPENDIXES

Appendix A	Summary of the Incidence of Neoplasms in Rats Administered Piperonyl Butoxide in the Diet	49
Table A1	Summary of the Incidence of Neoplasms in Male Rats Administered Piperonyl Butoxide in the Diet	51

		<u>Page</u>
Table A2	Summary of the Incidence of Neoplasms in Female Rats Administered Piperonyl Butoxide in the Diet	55
Appendix B	Summary of the Incidence of Neoplasms in Mice Administered Piperonyl Butoxide in the Diet	59
Table B1	Summary of the Incidence of Neoplasms in Male Mice Administered Piperonyl Butoxide in the Diet	61
Table B2	Summary of the Incidence of Neoplasms in Female Mice Administered Piperonyl Butoxide in the Diet	65
Appendix C	Summary of the Incidence of Nonneoplastic Lesions in Rats Administered Piperonyl Butoxide in the Diet	69
Table C1	Summary of the Incidence of Nonneoplastic Lesions in Male Rats Administered Piperonyl Butoxide in the Diet	71
Table C2	Summary of the Incidence of Nonneoplastic Lesions in Female Rats Administered Piperonyl Butoxide in the Diet	75
Appendix D	Summary of the Incidence of Nonneoplastic Lesions in Mice Administered Piperonyl Butoxide in the Diet	79
Table D1	Summary of the Incidence of Nonneoplastic Lesions in Male Mice Administered Piperonyl Butoxide in the Diet	81
Table D2	Summary of the Incidence of Nonneoplastic Lesions in Female Mice Administered Piperonyl Butoxide in the Diet	85
Appendix E	Analyses of the Incidence of Primary Tumors in Rats Administered Piperonyl Butoxide in the Diet	89

		<u>Page</u>
Table E1	Analyses of the Incidence of Primary Tumors in Male Rats Administered Piperonyl Butoxide in the Diet	91
Table E2	Analyses of the Incidence of Primary Tumors in Female Rats Administered Piperonyl Butoxide in the Diet	97
Appendix F	Analyses of the Incidence of Primary Tumors in Mice Administered Piperonyl Butoxide in the Diet	101
Table F1	Analyses of the Incidence of Primary Tumors in Male Mice Administered Piperonyl Butoxide in the Diet	103
Table F2	Analyses of the Incidence of Primary Tumors in Female Mice Administered Piperonyl Butoxide in the Diet	108

TABLES


Table 1	Piperonyl Butoxide Subchronic Feeding Studies in Rats and Mice	9
Table 2	Piperonyl Butoxide Chronic Feeding Studies in Rats	11
Table 3	Piperonyl Butoxide Chronic Feeding Studies in Mice	12

FIGURES

Figure 1	Growth Curves for Rats Administered Piperonyl Butoxide in the Diet	22
Figure 2	Survival Curves for Rats Administered Piperonyl Butoxide in the Diet	23
Figure 3	Growth Curves for Mice Administered Piperonyl Butoxide in the Diet	34
Figure 4	Survival Curves for Mice Administered Piperonyl Butoxide in the Diet	35

1	Introduction	1
2	Chapter 1: The History of the Book	2
3	Chapter 2: The Book in the Middle Ages	3
4	Chapter 3: The Book in the Renaissance	4
5	Chapter 4: The Book in the 17th and 18th Centuries	5
6	Chapter 5: The Book in the 19th Century	6
7	Chapter 6: The Book in the 20th Century	7
8	Chapter 7: The Book in the 21st Century	8
9	Chapter 8: The Book in the Future	9
10	Chapter 9: The Book in the Present	10
11	Chapter 10: The Book in the Past	11
12	Chapter 11: The Book in the Future	12
13	Chapter 12: The Book in the Present	13
14	Chapter 13: The Book in the Past	14
15	Chapter 14: The Book in the Future	15
16	Chapter 15: The Book in the Present	16
17	Chapter 16: The Book in the Past	17
18	Chapter 17: The Book in the Future	18
19	Chapter 18: The Book in the Present	19
20	Chapter 19: The Book in the Past	20
21	Chapter 20: The Book in the Future	21
22	Chapter 21: The Book in the Present	22
23	Chapter 22: The Book in the Past	23
24	Chapter 23: The Book in the Future	24
25	Chapter 24: The Book in the Present	25
26	Chapter 25: The Book in the Past	26
27	Chapter 26: The Book in the Future	27
28	Chapter 27: The Book in the Present	28
29	Chapter 28: The Book in the Past	29
30	Chapter 29: The Book in the Future	30
31	Chapter 30: The Book in the Present	31
32	Chapter 31: The Book in the Past	32
33	Chapter 32: The Book in the Future	33
34	Chapter 33: The Book in the Present	34
35	Chapter 34: The Book in the Past	35
36	Chapter 35: The Book in the Future	36
37	Chapter 36: The Book in the Present	37
38	Chapter 37: The Book in the Past	38
39	Chapter 38: The Book in the Future	39
40	Chapter 39: The Book in the Present	40
41	Chapter 40: The Book in the Past	41
42	Chapter 41: The Book in the Future	42
43	Chapter 42: The Book in the Present	43
44	Chapter 43: The Book in the Past	44
45	Chapter 44: The Book in the Future	45
46	Chapter 45: The Book in the Present	46
47	Chapter 46: The Book in the Past	47
48	Chapter 47: The Book in the Future	48
49	Chapter 48: The Book in the Present	49
50	Chapter 49: The Book in the Past	50
51	Chapter 50: The Book in the Future	51
52	Chapter 51: The Book in the Present	52
53	Chapter 52: The Book in the Past	53
54	Chapter 53: The Book in the Future	54
55	Chapter 54: The Book in the Present	55
56	Chapter 55: The Book in the Past	56
57	Chapter 56: The Book in the Future	57
58	Chapter 57: The Book in the Present	58
59	Chapter 58: The Book in the Past	59
60	Chapter 59: The Book in the Future	60
61	Chapter 60: The Book in the Present	61
62	Chapter 61: The Book in the Past	62
63	Chapter 62: The Book in the Future	63
64	Chapter 63: The Book in the Present	64
65	Chapter 64: The Book in the Past	65
66	Chapter 65: The Book in the Future	66
67	Chapter 66: The Book in the Present	67
68	Chapter 67: The Book in the Past	68
69	Chapter 68: The Book in the Future	69
70	Chapter 69: The Book in the Present	70
71	Chapter 70: The Book in the Past	71
72	Chapter 71: The Book in the Future	72
73	Chapter 72: The Book in the Present	73
74	Chapter 73: The Book in the Past	74
75	Chapter 74: The Book in the Future	75
76	Chapter 75: The Book in the Present	76
77	Chapter 76: The Book in the Past	77
78	Chapter 77: The Book in the Future	78
79	Chapter 78: The Book in the Present	79
80	Chapter 79: The Book in the Past	80
81	Chapter 80: The Book in the Future	81
82	Chapter 81: The Book in the Present	82
83	Chapter 82: The Book in the Past	83
84	Chapter 83: The Book in the Future	84
85	Chapter 84: The Book in the Present	85
86	Chapter 85: The Book in the Past	86
87	Chapter 86: The Book in the Future	87
88	Chapter 87: The Book in the Present	88
89	Chapter 88: The Book in the Past	89
90	Chapter 89: The Book in the Future	90
91	Chapter 90: The Book in the Present	91
92	Chapter 91: The Book in the Past	92
93	Chapter 92: The Book in the Future	93
94	Chapter 93: The Book in the Present	94
95	Chapter 94: The Book in the Past	95
96	Chapter 95: The Book in the Future	96
97	Chapter 96: The Book in the Present	97
98	Chapter 97: The Book in the Past	98
99	Chapter 98: The Book in the Future	99
100	Chapter 99: The Book in the Present	100
101	Chapter 100: The Book in the Past	101
102	Chapter 101: The Book in the Future	102
103	Chapter 102: The Book in the Present	103
104	Chapter 103: The Book in the Past	104
105	Chapter 104: The Book in the Future	105
106	Chapter 105: The Book in the Present	106
107	Chapter 106: The Book in the Past	107
108	Chapter 107: The Book in the Future	108
109	Chapter 108: The Book in the Present	109
110	Chapter 109: The Book in the Past	110
111	Chapter 110: The Book in the Future	111
112	Chapter 111: The Book in the Present	112
113	Chapter 112: The Book in the Past	113
114	Chapter 113: The Book in the Future	114
115	Chapter 114: The Book in the Present	115
116	Chapter 115: The Book in the Past	116
117	Chapter 116: The Book in the Future	117
118	Chapter 117: The Book in the Present	118
119	Chapter 118: The Book in the Past	119
120	Chapter 119: The Book in the Future	120
121	Chapter 120: The Book in the Present	121
122	Chapter 121: The Book in the Past	122
123	Chapter 122: The Book in the Future	123
124	Chapter 123: The Book in the Present	124
125	Chapter 124: The Book in the Past	125
126	Chapter 125: The Book in the Future	126
127	Chapter 126: The Book in the Present	127
128	Chapter 127: The Book in the Past	128
129	Chapter 128: The Book in the Future	129
130	Chapter 129: The Book in the Present	130
131	Chapter 130: The Book in the Past	131
132	Chapter 131: The Book in the Future	132
133	Chapter 132: The Book in the Present	133
134	Chapter 133: The Book in the Past	134
135	Chapter 134: The Book in the Future	135
136	Chapter 135: The Book in the Present	136
137	Chapter 136: The Book in the Past	137
138	Chapter 137: The Book in the Future	138
139	Chapter 138: The Book in the Present	139
140	Chapter 139: The Book in the Past	140
141	Chapter 140: The Book in the Future	141
142	Chapter 141: The Book in the Present	142
143	Chapter 142: The Book in the Past	143
144	Chapter 143: The Book in the Future	144
145	Chapter 144: The Book in the Present	145
146	Chapter 145: The Book in the Past	146
147	Chapter 146: The Book in the Future	147
148	Chapter 147: The Book in the Present	148
149	Chapter 148: The Book in the Past	149
150	Chapter 149: The Book in the Future	150
151	Chapter 150: The Book in the Present	151
152	Chapter 151: The Book in the Past	152
153	Chapter 152: The Book in the Future	153
154	Chapter 153: The Book in the Present	154
155	Chapter 154: The Book in the Past	155
156	Chapter 155: The Book in the Future	156
157	Chapter 156: The Book in the Present	157
158	Chapter 157: The Book in the Past	158
159	Chapter 158: The Book in the Future	159
160	Chapter 159: The Book in the Present	160
161	Chapter 160: The Book in the Past	161
162	Chapter 161: The Book in the Future	162
163	Chapter 162: The Book in the Present	163
164	Chapter 163: The Book in the Past	164
165	Chapter 164: The Book in the Future	165
166	Chapter 165: The Book in the Present	166
167	Chapter 166: The Book in the Past	167
168	Chapter 167: The Book in the Future	168
169	Chapter 168: The Book in the Present	169
170	Chapter 169: The Book in the Past	170
171	Chapter 170: The Book in the Future	171
172	Chapter 171: The Book in the Present	172
173	Chapter 172: The Book in the Past	173
174	Chapter 173: The Book in the Future	174
175	Chapter 174: The Book in the Present	175
176	Chapter 175: The Book in the Past	176
177	Chapter 176: The Book in the Future	177
178	Chapter 177: The Book in the Present	178
179	Chapter 178: The Book in the Past	179
180	Chapter 179: The Book in the Future	180
181	Chapter 180: The Book in the Present	181
182	Chapter 181: The Book in the Past	182
183	Chapter 182: The Book in the Future	183
184	Chapter 183: The Book in the Present	184
185	Chapter 184: The Book in the Past	185
186	Chapter 185: The Book in the Future	186
187	Chapter 186: The Book in the Present	187
188	Chapter 187: The Book in the Past	188
189	Chapter 188: The Book in the Future	189
190	Chapter 189: The Book in the Present	190
191	Chapter 190: The Book in the Past	191
192	Chapter 191: The Book in the Future	192
193	Chapter 192: The Book in the Present	193
194	Chapter 193: The Book in the Past	194
195	Chapter 194: The Book in the Future	195
196	Chapter 195: The Book in the Present	196
197	Chapter 196: The Book in the Past	197
198	Chapter 197: The Book in the Future	198
199	Chapter 198: The Book in the Present	199
200	Chapter 199: The Book in the Past	200
201	Chapter 200: The Book in the Future	201
202	Chapter 201: The Book in the Present	202
203	Chapter 202: The Book in the Past	203
204	Chapter 203: The Book in the Future	204
205	Chapter 204: The Book in the Present	205
206	Chapter 205: The Book in the Past	206
207	Chapter 206: The Book in the Future	207
208	Chapter 207: The Book in the Present	208
209	Chapter 208: The Book in the Past	209
210	Chapter 209: The Book in the Future	210
211	Chapter 210: The Book in the Present	211
212	Chapter 211: The Book in the Past	212
213	Chapter 212: The Book in the Future	213
214	Chapter 213: The Book in the Present	214
215	Chapter 214: The Book in the Past	215
216	Chapter 215: The Book in the Future	216
217	Chapter 216: The Book in the Present	217
218	Chapter 217: The Book in the Past	218
219	Chapter 218: The Book in the Future	219
220	Chapter 219: The Book in the Present	220
221	Chapter 220: The Book in the Past	221
222	Chapter 221: The Book in the Future	222
223	Chapter 222: The Book in the Present	223
224	Chapter 223: The Book in the Past	224
225	Chapter 224: The Book in the Future	225
226	Chapter 225: The Book in the Present	226
227	Chapter 226: The Book in the Past	227
228	Chapter 227: The Book in the Future	228
229	Chapter 228: The Book in the Present	229
230	Chapter 229: The Book in the Past	230
231	Chapter 230: The Book in the Future	231
232	Chapter 231: The Book in the Present	232
233	Chapter 232: The Book in the Past	233
234	Chapter 233: The Book in the Future	234
235	Chapter 234: The Book in the Present	235
236	Chapter 235: The Book in the Past	236
237	Chapter 236: The Book in the Future	237
238	Chapter 237: The Book in the Present	238
239	Chapter 238: The Book in the Past	239
240	Chapter 239: The Book in the Future	240
241	Chapter 240: The Book in the Present	241
242	Chapter 241: The Book in the Past	242
243	Chapter 242: The Book in the Future	243
244	Chapter 243: The Book in the Present	244
245	Chapter 244: The Book in the Past	245
246	Chapter 245: The Book in the Future	246
247	Chapter 246: The Book in the Present	247
248	Chapter 247: The Book in the Past	248
249	Chapter 248: The Book in the Future	249
250	Chapter 249: The Book in the Present	250
251	Chapter 250: The Book in the Past	251
252	Chapter 251: The Book in the Future	252
253	Chapter 252: The Book in the Present	253
254	Chapter 253: The Book in the Past	254
255	Chapter 254: The Book in the Future	255
256	Chapter 255: The Book in the Present	256
257	Chapter 256: The Book in the Past	257
258	Chapter 257: The Book in the Future	258
259	Chapter 258: The Book in the Present	259
260	Chapter 259: The Book in the Past	260
261	Chapter 260: The Book in the Future	261
262	Chapter 261: The Book in the Present	262
263	Chapter 262: The Book in the Past	263
264	Chapter 263: The Book in the Future	264
265	Chapter 264: The Book in the Present	265
266	Chapter 265: The Book in the Past	266
267	Chapter 266: The Book in the Future	267
268	Chapter 267: The Book in the Present	268
269	Chapter 268: The Book in the Past	269
270	Chapter 269: The Book in the Future	270
271	Chapter 270: The Book in the Present	271
272	Chapter 271: The Book in the Past	272
273	Chapter 272: The Book in the Future	273
274	Chapter 273: The Book in the Present	274
275	Chapter 274: The Book in the Past	275
276	Chapter 275: The Book in the Future	276
277	Chapter 276: The Book in the Present	277
278	Chapter 277: The Book in the Past	278
279	Chapter 278: The Book in the Future	279
280	Chapter 279: The Book in the Present	280
281	Chapter 280: The Book in the Past	281
282	Chapter 281: The Book in the Future	282
283	Chapter 282: The Book in the Present	283
284	Chapter 283: The Book in the Past	284
285	Chapter 284: The Book in the Future	285
286	Chapter 285: The Book in the Present	286
287	Chapter 286: The Book in the Past	287
288	Chapter 287: The Book in the Future	288
289	Chapter 288: The Book in the Present	289
290	Chapter 289: The Book in the Past	290
291	Chapter 290: The Book in the Future	291
292	Chapter 291: The Book in the Present	292
293	Chapter 292: The Book in the Past	293
294	Chapter 293: The Book in the Future	294
295	Chapter 294: The Book in the Present	295
296	Chapter 295: The Book in the Past	296
297	Chapter 296: The Book in the Future	297
298	Chapter 297: The Book in the Present	298
299	Chapter 298: The Book in the Past	299
300	Chapter 299: The Book in the Future	300
301	Chapter 300: The Book in the Present	301
302	Chapter 301	

I. INTRODUCTION



Piperonyl butoxide

Piperonyl butoxide (CAS 51-03-6; NCI C02813), 5-((2-(2-butoxyethoxy)ethoxy)-methyl-6-propyl-1,3-benzodioxole), is used to enhance the insecticidal properties of the pyrethrins, by blocking the pyrethrin detoxification enzymes in the insect (Metcalf, 1966). Pyrethrins alone produce a very rapid knockdown of insects, followed by substantial recovery, whereas addition of a synergist such as piperonyl butoxide decreases the insecticidal dose of pyrethrin (Metcalf, 1966). Piperonyl butoxide is also formulated with synthetic pyrethrin analogues, such as allethrin and tetramethrin (Stanford Research Institute, 1976).

Piperonyl butoxide has commercial importance as a synergist for

insecticides. Approximately 1 million pounds of this chemical were used in the United States in 1974, of which 80% was used in commercial, domestic (house and garden), and industrial establishments, and 20% on livestock and poultry (Ayers and Johnson, 1976).

The acute oral LD₅₀ of piperonyl butoxide has been reported as 11.5 ml/kg for rats of unspecified strain (Draize et al., 1948) and as between 7.5 and 10.0 gm/kg for Wistar rats (Sarles et al., 1949); that for mice (strains not specified) has been reported as 8.3 ml/kg (Draize et al., 1948) and as 3,800 mg/kg (Kenaga and Allison, 1969).

The long-term toxicity of piperonyl butoxide was investigated by Innes et al. (1969) as a part of a large-scale test of industrial and agricultural chemicals. These investigators did not obtain a significant incidence of tumors in either the (C57BL/6 x C₃H/Anf)F1 or the (C57BL/6 x AKR)F1 hybrid mice, the only species tested, and they categorized this chemical among those that probably required further testing. On the basis of these preliminary results, piperonyl butoxide was selected for study in the Carcinogenesis Testing Program.

II. MATERIALS AND METHODS

A. Chemical

The test chemical used in the bioassay was obtained as a 91-kilogram batch of Lot No. 5 technical-grade piperonyl butoxide from Niagara Chemical Company, FMC Corporation, Middleport, New York. Gas chromatographic (gc) analysis performed at Frederick Cancer Research Center, Frederick, Maryland, showed three major components and some minor ones totaling less than 1% each. As identified by the manufacturer, the three major components and their respective percentages of the total gc peak area were piperonyl butoxide, 88.4%; butyl carbitol, 2.1%; and an isomer of piperonyl butoxide, 2.4%. The infrared absorption spectra were identical with that of an authentic standard. The technical-grade piperonyl butoxide as described above will hereinafter be referred to in this report as piperonyl butoxide.

The piperonyl butoxide was stored at 7°C until used.

B. Dietary Preparation

Test diets containing piperonyl butoxide were prepared fresh each week in 6- to 12-kilogram batches at appropriate doses. A known weight of the chemical was first mixed with an equal weight of autoclaved Wayne® Sterilizable Lab Meal (Allied Mills, Inc., Chicago, Ill.), using a mortar and pestle. The mixing was repeated with second and third additions of feed, and final mixing was performed with the remaining quantity of feed for a minimum of 15 minutes in a Patterson-Kelly twin-shell blender.

The diets were stored at 7°C in plastic bags during the 1 to 1-1/2 weeks it was used.

C. Animals

Male and female Fischer 344 rats and B6C3F1 mice were obtained from the Frederick Cancer Research Center (Frederick, Md.) as 4-week-old weanlings, all within 3 days of the same age. The animals were housed within the test facility for 2 weeks and then were assigned four rats to a cage and five mice to a cage by a system that averaged the weights per cage for a given species and sex. For use in the chronic study, the male rats were required

to weigh 90 to 105 g, averaging at least 100 g; the female rats, 80 to 95 g, averaging at least 90 g; the male mice, 18 to 22 g, averaging at least 19.5 g; and the female mice, 17 to 21 g, averaging at least 18.5 g. Individual animals were identified by ear punch.

D. Animal Maintenance

The animals were housed in polycarbonate cages (Lab Products, Inc., Garfield, N.J.), 19 x 10-1/2 x 8 inches for the rats and 11-1/2 x 7-1/2 x 5 inches for the mice. The cages were suspended from aluminum racks (Scientific Cages, Inc., Bryan, Tex.) and were covered by nonwoven polyester-fiber 12-mil-thick filter paper (Hoeltge, Inc., Cincinnati, Ohio). The bedding used was Absorb-dri® hardwood chips (Northeastern Products, Inc., Warrenburg, N.Y.). The feed supplied was autoclaved Wayne® Sterilizable Lab Meal with 4% fat, provided ad libitum in suspended stainless steel hoppers and replenished at least three times per week. Water, acidified to pH 2.5, was supplied ad libitum from glass bottles and sipper tubes (Lab Products, Inc.) suspended through the tops of the cages.

Contaminated bedding was disposed of through an enclosed vacuum

line that led to a holding tank from which the bedding was fed periodically into an incinerator. The cages were sanitized at 82-88°C in a tunnel-type cagewasher (Industrial Washing Corp., Mataway, N. J.) twice per week, using the detergents, Clout® (Pharmaceutical Research Laboratories, Greenwich, Conn.) or Oxford D'Chlor (Oxford Chemicals, Atlanta, Ga.). The feed hoppers were sanitized twice per month in the same equipment. The glass bottles and sipper tubes were sanitized at 82-88°C in a tunnel-type bottle washer (Consolidated Equipment Supply Co., Mercersburg, Pa.) three times per week, using a Calgen Commercial Division detergent (St. Louis, Mo.). The racks for the cages were sanitized at or above 82°C in a rack washer (Consolidated Equipment Supply Co.) once per month, using the Calgen Commercial Division detergent, and the filter paper was changed at the same time.

The air in the animal rooms was regulated automatically at a temperature of 22-24°C and a relative humidity of 45-55%. Nonrecirculated air was passed through a filter of 65% efficiency and a bag filter of 95% efficiency at the intake and through a "Z"-type roughing filter of 30% efficiency and a bag system of 90-95% efficiency at the exhaust (American Air Filters, Louisville, Ky.; Mine Safety Appliances, Pittsburgh, Pa.). The rate of movement allowed 15 changes of room air per hour. The

air pressure was maintained negative to a clean hallway and positive to a return hallway. Fluorescent lighting was provided on a 12-hour-per-day cycle.

All control and dosed rats were housed in the same room as rats on feeding studies of the following chemicals:

(CAS 999-81-5) chlorocholine chloride
(CAS 88-06-2) 2,4,6-trichlorophenol

All control and dosed mice were housed in the same room as mice on feeding studies of the following chemicals:

(CAS 128-66-5) C. I. vat yellow 4 (amanthrene)
(CAS 103-33-3) azobenzene
(CAS 20941-65-5) ethyl tellurac
(CAS 88-06-2) 2,4,6-trichlorophenol
(CAS 72-56-0) p,p'-ethyl-DDD
(CAS 85-44-9) phthalic anhydride

E. Subchronic Studies

Subchronic feeding studies were conducted to estimate the maximum tolerated doses (MTDs) of piperonyl butoxide, on the basis of which two concentrations (hereinafter referred to as "low" and "high" doses) were selected for administration in the chronic studies. Groups of five rats of each sex and five mice of each sex were administered feed containing piperonyl butoxide at one

of several doses, and groups of five control animals of each species and sex were administered a basal diet only. The doses used for the rats in the subchronic studies were 14,700, 21,500, 31,500, 46,000, and 68,000 ppm; the doses used for the mice in the subchronic studies were 4,600, 6,800, 10,000, 14,700, 21,500, 31,500, 46,000, and 68,000 ppm. The period of administration of the test chemical was 7 weeks, followed by 1 week of additional observation. Each animal was weighed twice per week. Table 1 shows the number of animals dying at each dose and the number of weeks on study when the deaths occurred; the table also shows the mean body weights of dosed animals at week 7, expressed as percentages of mean body weights of controls at week 7.

At the end of the subchronic studies, all animals were killed by CO₂ inhalation and necropsied. The lowest doses at which clinical and histopathologic findings were observed were 21,500 ppm in the male and female rats and 14,700 ppm in the male and female mice. In the rats, interstitial pneumonia, perivascular lymphoplasmacytosis, and peribronchiolar lymphocytosis were observed, but were not considered to be related to administration of the test chemical. In the mice, extramedullary hematopoiesis was observed in one male and five females, polypoid nuclei were observed in livers of two males and one female, and slight vacuolization of hepatocytes was observed in four females.

Table 1

Piperonyl Butoxide Subchronic Feeding Studies in Rats and Mice

(ppm)	Male			Female		
	Mortality		Mean Weight at Week 7 as % of Control	Mortality		Mean Weight at Week 7 as % of Control
	Number Dead	Week on Study		Number Dead	Week on Study	
<u>RATS</u>						
14,700			94			95
21,500			73			76
31,500			60			68
46,000	4	8	60	3	8	38
68,000	5	1		5	1	
<u>MICE</u>						
4,600			91			94
6,800			87			93
10,000			76	1	8	76
14,700	1	8	85			72
21,500	4	8	76	4	8	87
31,000	5	2		4	8	
46,000	5	2		5	2	
68,000	5	2		5	2	

Ten percent depression in body weight was taken as the major criterion for estimation of MTDs. The doses required to produce this response were determined by the following procedure: first, least squares regressions of mean body weights versus days on study were used to estimate mean body weights of each of the dosed groups at day 49. Next, probits of the percent weights of dosed groups at day 49 relative to weights of corresponding control groups were plotted against logarithms of the doses, and least squares regressions fitted to the data were used to estimate the doses required to induce 10% depression in weight.

When these data were evaluated in conjunction with the clinical and histopathologic findings, the low and high doses for chronic studies using rats were set at 5,000 and 10,000 ppm, and those for chronic studies using mice were set at 2,500 and 5,000 ppm.

F. Chronic Studies

The test groups, doses administered, and durations of the chronic feeding studies are shown in tables 2 and 3. Doses administered to the mice were reduced after week 30 as indicated.

Table 2. Piperonyl Butoxide Chronic Feeding Studies in Rats

<u>Sex and Test Group</u>	<u>Initial No. of Animals (a)</u>	<u>Piperonyl Butoxide in Diet (b) (ppm)</u>	<u>Time on Study (weeks)</u>
<u>Male</u>			
Matched-Control	20	0	107
Low-Dose	50	5,000	107
High-Dose	50	10,000	107
<u>Female</u>			
Matched-Control	20	0	107
Low-Dose	50	5,000	107
High-Dose	50	10,000	107

(a) All animals were 6 weeks of age when placed on study.

(b) Test and control diets were provided ad libitum.

Table 3. Piperonyl Butoxide Chronic Feeding Studies in Mice

<u>Sex and Test Group</u>	<u>Initial No. of Animals (a)</u>	<u>Piperonyl Butoxide in Diet (b) (ppm)</u>	<u>Time on Study (weeks)</u>	<u>Time-Weighted Average Dose (c) (ppm)</u>
<u>Male</u>				
Matched-Control	20	0	112	
Low-Dose	50	2,500	30	
		500	82	1,036
High-Dose	50	5,000	30	
		2,000	82	2,804
<u>Female</u>				
Matched-Control	20	0	112	
Low-Dose	50	2,500	30	
		500	82	1,036
High-Dose	50	5,000	30	
		2,000	82	2,804

(a) All animals were 6 weeks of age when placed on study.

(b) Test and control diets were provided ad libitum.

(c) Time-weighted average dose =
$$\frac{\Sigma(\text{dose (ppm)} \times \text{no. of weeks at that dose})}{\Sigma(\text{no. of weeks receiving each dose})}$$

G. Clinical and Pathologic Examinations

All animals were observed twice daily for signs of toxicity. Clinical examination and palpation for masses were performed each month, and the animals were weighed at least once per month. Moribund animals and animals that survived to the end of the bioassay were killed by asphyxiation using CO₂ and necropsied. Necropsies were also performed on all animals found dead, unless precluded by autolysis or severe cannibalization.

The pathologic evaluation consisted of gross and microscopic examination of major tissues, major organs, and all gross lesions. The tissues were preserved in 10% buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin. The following tissues were examined microscopically: skin, lungs and bronchi, trachea, bone marrow (femur), spleen, lymph nodes (mesenteric and submandibular), thymus, heart, salivary glands (parotid, sublingual, and submaxillary), liver, pancreas, esophagus, stomach (glandular and nonglandular), small and large intestine, kidney, urinary bladder, pituitary, adrenal, thyroid, parathyroid, testis, prostate, mammary gland, uterus, ovary, brain (cerebrum and cerebellum), and all tissue masses. Peripheral blood smears also were made for all animals, whenever possible.

A few tissues from some animals were not examined, particularly from those animals that may have died early, been missing, or been in advanced states of cannibalization or autolysis. Thus, the number of animals from which particular organs or tissues were examined microscopically varies, and does not necessarily represent the number of animals that were placed on study in each group.

H. Data Recording and Statistical Analyses

Pertinent data on this experiment have been recorded in an automatic data processing system, the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969). Data tables were generated for verification of data transcription and for statistical review. These data were analyzed using the appropriate statistical techniques described in this section. Those analyses of the experimental results that bear on the possibility of carcinogenicity are discussed in the statistical narrative section.

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) for testing two groups for equality and Tarone's (1975) extensions of Cox's methods for testing for a dose-related trend. One-tailed P values have been reported for all tests except the departure from linearity test, which is only reported when its two-tailed P value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given the ratio as of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site is examined (denominator). In most instances, the denominators included only those animals for which that site was examined histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970) was used to compare the tumor incidence of a control group with that of a group of dosed animals at each dose level. When results for a number of dosed groups (k) are compared simultaneously with those for a control group, a correction to ensure an overall significance level of 0.05 may be made. The Bonferroni inequality (Miller, 1966) requires that the P value for any comparison be less than or equal to $0.05/k$. In cases where this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971), was also used. Under the assumption of a linear trend, this test determines if the slope of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend is a positive dose relationship. This method also provides a two-tailed test of departure from linear trend.

A time-adjusted analysis was applied when numerous early deaths resulted from causes that were not associated with the formation of tumors. In this analysis, deaths that occurred before the first tumor was observed were excluded by basing the statistical tests on animals that survived at least 52 weeks, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at least as long as the animal in which the first tumor was found. Once this reduced set of data was obtained, the standard procedures for analyses of the incidence of tumors (Fisher exact tests, Cochran-Armitage tests, etc.) were followed.

When appropriate, life-table methods were used to analyze the incidence of tumors. Curves of the proportions surviving without an observed tumor were computed as in Saffiotti et al. (1972). The week during which an animal died naturally or was sacrificed was entered as the time point of tumor observation. Cox's methods of comparing these curves were used for two groups; Tarone's extension to testing for linear trend was used for three groups. The statistical tests for the incidence of tumors which used life-table methods were one-tailed and, unless otherwise noted, in the direction of a positive dose relationship.

Significant departures from linearity (P less than 0.05, two-tailed test) were also noted.

The approximate 95 percent confidence interval for the relative risk of each dosed group compared with its control was calculated from the exact interval on the odds ratio (Gart, 1971). The relative risk is defined as p_t/p_c where p_t is the true binomial probability of the incidence of a specific type of tumor in a dosed group of animals and p_c is the true probability of the spontaneous incidence of the same type of tumor in a control group. The hypothesis of equality between the true proportion of a specific tumor in a dosed group and the proportion in a control group corresponds to a relative risk of unity. Values in excess of unity represent the condition of a larger proportion in the dosed group than in the control.

The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that in approximately 95% of a large number of identical experiments, the true ratio of the risk in a dosed group of animals to that in a control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically

significant result (P less than 0.025 one-tailed test when the control incidence is not zero, P less than 0.050 when the control incidence is zero) has occurred. When the lower limit is less than unity, but the upper limit is greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility of the induction of tumors by the test chemical, which could not be detected under the conditions of this test.

III. RESULTS - RATS

A. Body Weights and Clinical Signs (Rats)

A dose-related depression in the mean body weights of both dosed male and female rats, when compared with corresponding matched controls, was observed throughout the bioassay (figure 1). However, the depressions in the mean body weights of the low- and high-dose groups were slight during the first 50 weeks. Wasting among the rats and redness of the skin and mucous membranes occurred at low incidences in some of the dosed groups and may have been related to administration of the test chemical. Other clinical signs, including corneal opacity and tissue masses, occurred at comparable incidences in dosed and control groups.

B. Survival (Rats)

The Kaplan and Meier curves estimating the probabilities of survival for male and female rats administered piperonyl butoxide in the diet at the doses of this bioassay, together with those of the matched controls, are shown in figure 2. The result of the Tarone test for dose-related trend in mortality is not significant in either sex.





Figure 1. Growth Curves for Rats Administered Piperonyl Butoxide in the Diet





Figure 2. Survival Curves For Rats Administered Piperonyl Butoxide in the Diet

In male rats, 37/50 (74%) of the high-dose group, 36/50 (72%) of the low-dose group, and 15/20 (75%) of the controls were alive at termination of the study. In females, 40/50 (80%) of the high-dose group, 41/50 (82%) of the low-dose group, and all 20 of the controls were alive at termination of the study.

Sufficient numbers of rats of each sex were at risk for the development of late-appearing tumors.

C. Pathology (Rats)

Histopathologic findings on neoplasms in rats are summarized in Appendix A, tables A1 and A2; findings on nonneoplastic lesions are summarized in Appendix C, tables C1 and C2.

With regard to liver lesions, the term "focal hyperplasia" was used in this study to indicate the presence of one or more foci of hepatocytes with increased cytoplasmic basophilia and a slight increase in the amount of nuclear chromatin. Many of these hepatocytes also had a slight increase in nuclear:cytoplasmic ratio when compared with adjacent normal hepatocytes, and, infrequently, mitotic figures or hepatocytes with double nuclei were observed. These foci of hyperbasophilic hepatocytes were thought to represent areas of hyperplasia and were diagnosed as

such in this study. They did not compress adjacent hepatic parenchyma. These lesions are similar morphologically to those described by Squire and Levitt (1975) as "basophilic foci."

Lesions classified as "hepatocytomegaly" consisted of foci of enlarged hepatocytes, many of which contained large, vesicular nuclei and numerous fine cytoplasmic vacuoles, which gave the cytoplasm a "ground glass" appearance. Distortion of lobular architecture in these foci was minimal, and trabeculae were continuous with adjacent normal hepatocytes. These lesions correspond morphologically to those described by Squire and Levitt (1975) as "eosinophilic foci," "ground glass foci," or "clear cell foci."

Lesions classified as "neoplastic nodules" had many similarities to the previously described proliferative hepatocytic lesions. However, the lesions classified as neoplastic nodules were larger and contained more distinct abnormality of lobular architecture; liver cords at the periphery of the neoplastic nodules were oriented perpendicular to cords of adjacent normal hepatic parenchyma, and distinct compression of adjacent normal liver was evident.

Endocrine tissues were the most frequent sites of neoplasms in

both dosed and control rats in this study. Interstitial-cell tumors of the testis were observed in nearly all male rats in all groups; a high spontaneous incidence of this tumor is characteristic of aged Fischer 344 rats. Adenomas of the pituitary were also found at a high incidence in all groups, especially females. Other endocrine neoplasms observed included follicular-cell and C-cell tumors of the thyroid, islet-cell tumors of the pancreas, and pheochromocytomas and cortical carcinomas of the adrenal.

In some proliferative endocrine lesions, differentiation between benign and malignant neoplasms was difficult. C-cell lesions of the thyroid were classified as adenomas when the proliferating C cells were present in nodular masses that widely separated the thyroid follicles and distorted the follicular architecture. In some of the larger adenomas, the C cells were present in interlacing bundles of elongated, spindling cells, rather than the polyhedral to spherical shape characteristic of normal C cells. When invasion of thyroid capsule, adjacent tissues, or vessels was present, or when metastasis was detected, the lesion was classified as a C-cell carcinoma.

Follicular-cell neoplasms occurred less frequently than C-cell neoplasms. The follicular-cell adenomas appeared microscopically

as well-circumscribed masses composed of enlarged follicles lined by hyperbasophilic follicular cells which were increased in number per unit area by papillary infolding of simple cuboidal or columnar epithelium into the follicular lumen and by stratification of follicular cells surrounding the lumen. Distinct compression of adjacent normal thyroid parenchyma, with some evidence of fibrous encapsulation, was present. Follicular-cell lesions were classified as carcinomas based upon the presence of anaplasia and histologic arrangement in disorderly nests and/or sheets. Areas with papillary patterns were also present. Fibrous stroma often intermingled with, but did not encapsulate, follicular-cell carcinomas.

The diagnosis of pheochromocytoma was made when the adrenal medullary lesion was present as a discrete mass that compressed adjacent normal adrenal parenchyma. These neoplasms were composed of sheets, nests, and/or cords of polyhedral to spherical cells with abundant, slightly basophilic cytoplasm and large nuclei with abundant chromatin. Islet-cell adenomas appeared as discrete, encapsulated nodules of islet cells that compressed the adjacent normal pancreas; diagnosis of islet-cell carcinoma in a high-dose male rat was based on invasion of the capsule surrounding the neoplasm.

Two neoplasms were observed in the intestinal tract. A malignant spindle-cell tumor was present in the wall of the cecum of a high-dose male rat. This was a large neoplasm that extended into the mucosa; the diagnosis of leiomyosarcoma was based on the location in the wall and the cellular morphology. A hemangiosarcoma was present in the wall of the large intestine of a low-dose female rat.

Malignant lymphomas occurred rather frequently. Most of these neoplasms were composed of relatively undifferentiated lymphoreticular cells and involved numerous organs and tissues throughout the body. The organ most frequently affected was the spleen; also frequently affected were the liver, lymph nodes, thymus, and lungs. There was only one malignant lymphoma in the control group of female rats in this study; this is a relatively common tumor in Fischer 344 rats. Evidence of leukemia (masses of neoplastic lymphoreticular cells in vessel lumens) was seen in some cases.

Proliferative pulmonary lesions were observed rather infrequently in both dosed and control groups. Differentiation between adenomas and carcinomas was based on degree of anaplasia, mitotic index, size of the neoplasm, and presence of apparent invasion of adjacent pulmonary parenchyma in carcinomas, as opposed to mere

compression of adjacent parenchyma, and thus, a more discrete lesion in adenomas.

The most frequently occurring neoplasm of the reproductive tract, other than the previously mentioned interstitial-cell tumor of the testis, was the endometrial stromal polyp of the uterus. This lesion was present as a discrete mass protruding into the lumen of the uterus, lined by endometrium, and sometimes associated with suppurative endometritis and/or cystic endometrial hyperplasia. The stroma was usually proliferating in a rather loosely woven pattern, with numerous small vessels interspersed among stromal cells.

Mesothelioma of the tunica vaginalis was observed in sections of testis from four male rats, all of which were from dosed groups. One generalized peritoneal mesothelioma was present in a low-dose male rat.

Squamous-cell carcinomas were observed in the preputial gland of a high-dose male rat and in the clitoral gland of a high-dose female.

The most common neoplasms of the mammary gland were

fibroadenomas; these occurred only in females, were often multiple, and were seen in both dosed and control groups.

Various other types of malignant and benign neoplasms were observed at low incidences in sections of skin and subcutis, and in other organs and tissues throughout the body. No apparent difference in incidence of these neoplasms between dosed and control groups was present.

Numerous inflammatory, degenerative, and proliferative lesions commonly observed in aged Fischer 344 rats occurred with approximately equal frequency in dosed and control rats. These lesions included chronic tracheitis; multifocal alveolar macrophage aggregates in lung parenchyma; chronic nephritis with scarring, tubular dilatation, and tubular regeneration; ectasia of lymph sinuses in the mesenteric lymph nodes; testicular atrophy; and C-cell hyperplasia of the thyroid.

Other nonneoplastic proliferative lesions included hyperplasia of the follicular cells of the thyroid, adrenal medulla and cortex, parathyroid, endometrium, mammary epithelium, and hepatocytes.

Based on the histopathologic examination, there were instances in this study in which neoplastic or hyperplastic lesions occurred

only in dosed animals, or with increased frequency in dosed animals when compared with control groups. However, the nature, incidence, and severity of the lesions observed provide no clear evidence of carcinogenic effect of piperonyl butoxide in Fischer 344 rats under the conditions of this bioassay.

D. Statistical Analyses of Results (Rats)

Tables E1 and E2 in Appendix E contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals of one group and at an incidence of at least 5% in one or more than one group.

In female rats, the result of the Cochran-Armitage test for positive dose-related trend in the incidence of lymphomas is significant ($P = 0.007$). The results of the Fisher exact test show that the incidence in the high-dose group is significantly higher ($P = 0.020$) than that in the control group. The historical record of the incidence of lymphomas and leukemias in female rats at this laboratory to date is 19/191(10%). These historical-control groups include one with an incidence of animals with lymphoma or leukemia of 7/20 (35%) and another with an incidence of 6/20 (30%). The statistical conclusion suggests

that the incidence of lymphomas in female rats may be associated with the administration of piperonyl butoxide; however, this conclusion may be due to the lower than usual incidence in the control group compared with the historical data.

Significant results in the negative direction are observed in the incidences of neoplastic nodules of the liver and of adenomas or carcinomas of the pituitary in the male rats; the incidences of these tumors in the control group exceed those in the dosed groups.

In each of the 95% confidence intervals for relative risk shown in the tables, the value of one is included; this indicates the absence of significant positive results. It should also be noted that each of the intervals has an upper limit greater than one, indicating the theoretical possibility of the induction of tumors, which could not be detected under the conditions of this test.

IV. RESULTS - MICE

A. Body Weights and Clinical Signs (Mice)

A dose-related depression in the mean body weights of both dosed male and female mice, when compared with corresponding matched controls, was observed throughout the bioassay (figure 3). Alopecia occurred at low incidences in the dosed groups of females and may have been related to administration of the test chemical. Corneal opacity and tissue masses occurred at comparable incidences in dosed and control groups.

B. Survival (Mice)

The Kaplan and Meier curves estimating the probabilities of survival for male and female mice administered piperonyl butoxide in the diet at the doses of this bioassay, together with those of the matched controls, are shown in figure 4. The result of the Tarone test for dose-related trend in mortality is not significant in either sex.

In male mice, 41/50 (82%) of the high-dose group, 42/50 (84%) of





Figure 3. Growth Curves for Mice Administered Piperonyl Butoxide in the Diet





Figure 4. Survival Curves For Mice Administered Piperonyl Butoxide in the Diet

the low-dose group, and 17/20 (85%) of the control group survived to termination of the study. In females, 34/50 (68%) of the high-dose group, 35/50 (70%) of the low-dose group, and 15/20 (75%) of the control group lived to termination of the study.

Sufficient numbers of mice of each sex were at risk for the development of late-appearing tumors.

C. Pathology (Mice)

Histopathologic findings on neoplasms in mice are summarized in Appendix B, tables B1 and B2; findings on nonneoplastic lesions are summarized in Appendix D, tables D1 and D2.

The most common neoplasm in male mice was hepatocellular carcinoma. There were 10/20 in control males, 17/50 in low-dose males, and 20/50 in high-dose males. Metastases were found in two mice from each of these groups. There was one hepatocellular carcinoma in a control female, two in low-dose females, and five in high-dose females. Metastases to the lungs were found in one high-dose female.

The next most common tumors were of lymphoreticular origin

(malignant lymphoma and reticulum-cell sarcoma). Their incidences were as follows: males — controls 4/20, low-dose 11/49, high-dose 6/50; females — controls 8/20, low-dose 14/47, high-dose 9/50. Many of these tumors were widely disseminated, involving several organs and forming tissue masses.

One low-dose male had a widely disseminated round-cell tumor that was diagnosed as a mast-cell tumor.

Hemangiosarcomas were found in one control male, one control female, three low-dose males, two low-dose females, and in one male and one female in the high-dose groups. Both of the tumors in the low-dose females and the one in a high-dose male were in multiple organs.

There were occasional alveolar-cell adenomas and adenocarcinomas of the lung. They were distributed with approximately equal frequency between males and females and between dosed and control groups.

Most other proliferative or neoplastic lesions were of single occurrence or very low incidence with approximately equal distribution between the dosed and control groups. There were

four adenomas of the lacrimal gland in high-dose males; none were found in any of the female mice.

In addition to the proliferative lesions, there was a scattering of inflammatory and degenerative changes in some mice in each group. These included focal mineralization of the brain, myocarditis, nephritis, focal hepatic necrosis, pancreatic atrophy, cystic endometrium, and ovarian cysts.

Based on the histopathologic examination, there were no differences in the incidences of neoplastic or nonneoplastic lesions between dosed and control mice; it is concluded that the dietary administration of piperonyl butoxide was not carcinogenic in B6C3F1 mice under the conditions of this bioassay.

D. Statistical Analyses of Results (Mice)

Tables F1 and F2 in Appendix F contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals of one group and at an incidence of at least 5% in one or more than one group.

In male mice, the result of the Cochran-Armitage test for positive dose-related trend in the incidence of adenomas of the lacrimal gland is significant ($P = 0.022$), but the results of the Fisher exact test are not significant.

A significant dose-related trend in the negative direction is observed for the incidence of lymphomas in female mice; the incidence of this tumor in the control group exceeds the incidences in the dosed groups.

In each of the 95% confidence intervals of relative risk, shown in the tables, the value of one is included; this indicates the absence of significant positive results. It should also be noted that each of the intervals has an upper limit greater than one, indicating the theoretical possibility of the induction of tumors by piperonyl butoxide, which could not be detected under the conditions of this test.

The first part of the paper discusses the importance of the study of the history of the English language. It is argued that the study of the history of the English language is not only a matter of academic interest but also of practical importance. The study of the history of the English language can help us to understand the development of the English language and to see how it has changed over time. It can also help us to understand the relationship between the English language and other languages.

The second part of the paper discusses the importance of the study of the history of the English language. It is argued that the study of the history of the English language is not only a matter of academic interest but also of practical importance. The study of the history of the English language can help us to understand the development of the English language and to see how it has changed over time. It can also help us to understand the relationship between the English language and other languages.

The third part of the paper discusses the importance of the study of the history of the English language. It is argued that the study of the history of the English language is not only a matter of academic interest but also of practical importance. The study of the history of the English language can help us to understand the development of the English language and to see how it has changed over time. It can also help us to understand the relationship between the English language and other languages.

The fourth part of the paper discusses the importance of the study of the history of the English language. It is argued that the study of the history of the English language is not only a matter of academic interest but also of practical importance. The study of the history of the English language can help us to understand the development of the English language and to see how it has changed over time. It can also help us to understand the relationship between the English language and other languages.

V. DISCUSSION

Dose-related depressions in mean body weight occurred during most or all of the bioassay in both rats and mice administered piperonyl butoxide. Redness of the skin and mucous membranes and wasting among the rats and alopecia among the mice occurred at low incidences in some of the dosed groups and may have been related to administration of the test chemical. Survival of the rats and mice was unaffected by the piperonyl butoxide and was 80% or greater in all groups at week 90 of the bioassay; thus, sufficient numbers of dosed and control rats and mice of each sex were at risk for the development of late-appearing tumors.

In the female rats, lymphomas occurred at incidences that were dose-related ($P = 0.007$); in a direct comparison, the incidence of the tumor in the high-dose group was higher ($P = 0.020$) than that in the control group (controls 1/20, low-dose 7/50, high-dose 15/50). However, the incidence of lymphomas, leukemias, and reticuloses in historical-control female Fischer 344 rats at the same laboratory was 19/191 (10%). These historical-control groups include one with an incidence of animals with lymphoma or leukemia of 7/20 (35%) and another with incidence of 6/20 (30%). Thus, the incidence of lymphomas in the

control female rats of the present bioassay may have been abnormally low, and the occurrence of the higher incidence in the dosed groups cannot be clearly related to administration of the piperonyl butoxide.

In the male mice, adenomas of the eye or lacrimal gland occurred at incidences that were dose related ($P = 0.023$), but in direct comparisons the incidences in the individual dosed groups were not significantly higher than that in the control group (controls 0/20, low-dose 0/49, high-dose 4/50); thus, the occurrence of this tumor in the male mice was not clearly related to administration of the test chemical.

When piperonyl butoxide was administered in the diet for 2 years to Wistar albino rats at doses as high as 25,000 ppm (Sarles and Vandegrift, 1952), no evidence was found for carcinogenicity. When the chemical was administered at 464 mg/kg by stomach tube for 3 weeks, then in the diet at 1,112 ppm for 18 months, to hybrid mice (C57BL/6 x C3H/Anf and C57BL/6 x AKR), an elevated incidence of reticulum-cell sarcoma was observed ($P = 0.05$) but was not significant at the selected criterion of $P = 0.01$ (NTIS, 1968; Innes et al., 1969). No evidence of carcinogenicity was observed when neonatal Swiss mice were given subcutaneous injections of 0.1 ml of a 5% solution of the chemical in

redistilled tricaprylin at days 1 and 7 and 0.2 ml at days 14 and 21; however, when the chemical was injected subcutaneously into the neonatal mice in combination with Freon 112, hepatomas developed at a statistically significant incidence (Epstein et al., 1967). Certain structural congeners of piperonyl butoxide (safrole, isosafrole, and dihydrosafrole) have been reported carcinogenic for rats (Osborne-Mandel) and mice, (C57BL/6 x C₃H/Anf and C57BL/6 x AKR) inducing tumors of the liver, esophagus, or lung, depending on species and sex (Long et al., 1963; Hagan et al., 1965; Innes et al., 1969).

Piperonyl butoxide is used commercially with pyrethins. This bioassay, however, tests the carcinogenicity of technical-grade piperonyl butoxide alone, and no conclusions can be drawn from the data in this report as to the possible carcinogenic effects of the combination of the two chemicals.

It is concluded that under the conditions of this bioassay, there was no convincing evidence that piperonyl butoxide was carcinogenic for Fischer 344 rats or B6C3F1 mice.

VI. BIBLIOGRAPHY

Armitage, P., Statistical Methods in Medical Research, John Wiley & Sons, Inc., New York, 1971, pp. 362-365.

Ayers, J. H. and Johnson, O. H., Insecticides In: Chemical Economics Handbook, Stanford Research Institute, Menlo Park, Calif., 1976, sec. 573.30071.

Berenblum, I., ed., Carcinogenicity Testing: A Report of the Panel on Carcinogenicity of the Cancer Research Commission of the UICC, Vol. 2. International Union Against Cancer, Geneva, 1969.

Cox, D. R., Regression models and life tables. J. R. Statist. Soc. B 34:187-220, 1972.

Cox, D. R., Analysis of Binary Data, Methuen & Co., Ltd. London, 1970, pp. 48-52.

Draize, J. H., Alvarez, E., Whitesell, M. F., Woodard, G., Hagan, E. C., and Nelson, A. A., Toxicological investigations of compounds proposed for use as insect repellents. J. Pharmacol. Exp. Therap. 93:26-39, 1948.

Epstein, S. S., Joshi, S., Andrea, J., Clapp, P., Falk, H., and Mantel, N., Synergistic toxicity and carcinogenicity of 'Freons' and piperonyl butoxide. Nature 214:526-528, 1967.

Gart, J. J., The comparison of proportions: a review of significance tests, confidence limits and adjustments for stratification. Rev. Int. Stat. Inst. 39:148-169, 1971.

Hagan, E. C., Jenner, P. M., Jones, W. I., Fitzhugh, O. G., Long, E. L., Brouwer, J. G., and Webb, W. K., Toxic properties of compounds related to safrole. Toxicol. Appl. Pharmacol. 7:18-24, 1965.

Innes, J. R. M., Ulland, B. M., Valerio, M. G., Petrucelli, L., Fishbein, L., Hart, E. R., Pallotta, A. J., Bates, R. R., Falk, H. L., Gart, J. J., Klein, M., Mitchell, I. and Peters, J., Bioassay of pesticides and industrial chemicals for tumorigenicity in mice: a preliminary note. J. Natl Cancer Inst. 42:1101-1114, 1969.

Kaplan, E. L. and Meier, P., Nonparametric estimation from incomplete observations. J. Am. Statist. Assoc. 53:457-481, 1958.

Kenaga, E. E., and Allison, W. E., Commercial and experimental organic insecticides. Bull. Entomol. Soc. Amer. 15 (2):85-148, 1969.

Linhart, M. S., Cooper, J. A., Martin, R. L., Page, N. P., and Peters, J. A., Carcinogenesis bioassay data system. Comp. and Biomed. Res. 7:230-248, 1974.

Long, E. L., Nelson, A. A., Fitzhugh, O. G., and Hansen, W. H., Liver tumors produced in rats by feeding safrole. Arch. Path. 75:595-604, 1963.

Metcalf, R. L., Insecticides. In: Kirk-Othmer Encyclopedia of Chemical Technology, Vol. 11, Interscience Publishers, New York, 1966, pp. 685-686.

Miller, R. G., Jr., Simultaneous Statistical Inference, McGraw-Hill Book Co., New York, 1966, pp. 6-10.

National Technical Information Service, Evaluation of Carcinogenic, Teratogenic, and Mutagenic Activities of Selected Pesticide and Industrial Chemicals, Vol. I, Carcinogenic Study, U. S. Department of Commerce. PB-223 159. August, 1968, p. 64.

Saffiotti, U., Montesano, R., Sellakumar, A. R., Cefis, F., and Kaufman, D. G., Respiratory tract carcinogenesis in hamsters induced by different numbers of administrations of benzo(a) pyrene and ferric oxide. Cancer Res. 32:1073-1081, 1972.

Sarles, M. P. and Vandegrift, W. B., Chronic oral toxicity and related studies on animals with the insecticide and pyrethrum synergist, piperonyl butoxide. Amer. J. Trop. Med. Hyg. 1:862-883, 1952.

Sarles, M. P., Dove, W. E., and Moore, D. H., Acute toxicity and irritation tests on animals with the new insecticide piperonyl butoxide. Am. J. Trop. Med. 29:151-166, 1949.

Squire, R. A. and Levitt, M. H., Report of a workshop on classification of specific hepatocellular lesions in rats. Cancer Res. 35:3214-3223, 1975.

Stanford Research Institute, Stage I Chemical Dossier. Stanford Research Institute, Menlo Park, Calif., April 1976.

Tarone, R. E., Tests for trend in life table analysis. Biometrika 62:679-682, 1975.

...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...

APPENDIX A

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN RATS ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1. Introduction

The following table shows the results of the experiment. The data is presented in a clear and concise manner, allowing for easy comparison of the different groups. The results are as follows:

TABLE A1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
KERATOACANTHOMA	1 (5%)	1 (2%)	
*SUBCUT TISSUE	(20)	(50)	(50)
FIBROMA		2 (4%)	3 (6%)
FIBROSARCOMA		1 (2%)	1 (2%)
LIPOMA		1 (2%)	
LIPOSARCOMA			1 (2%)
NEUPILEOMA, MALIGNANT			1 (2%)
RESPIRATORY SYSTEM			
#LUNG	(20)	(48)	(50)
ALVEOLAR/BRONCHIOLAR ADENOMA			2 (4%)
ALVEOLAR/BRONCHIOLAR CARCINOMA	1 (5%)	1 (2%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(50)
MALIGNANT LYMPHOMA, NOS	2 (10%)	3 (6%)	3 (6%)
MALIG.LYMPHOMA, UNDIFFER-TYPE	7 (35%)	12 (24%)	9 (18%)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE			1 (2%)
CIRCULATORY SYSTEM			
#HEART	(20)	(50)	(50)
ALVEOLAR/BRONCHIOLAR CA, INVASIV	1 (5%)		
DIGESTIVE SYSTEM			
*LIVER	(20)	(50)	(49)
NEOPLASTIC NODULE	2 (10%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
# CECUM LEIOMYOSARCOMA	(20)	(49)	(48) 1 (2%)
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
# PITUITARY	(19)	(49)	(48)
CARCINOMA, NOS			1 (2%)
ADENOMA, NOS	4 (21%)		
CHROMOPHOBE ADENOMA	1 (5%)	7 (14%)	6 (13%)
# ADRENAL	(20)	(50)	(50)
CORTICAL ADENOMA	1 (5%)		
PHEOCHROMOCYTOMA	2 (10%)	4 (8%)	4 (8%)
# THYROID	(20)	(49)	(50)
FOLLICULAR-CELL ADENOMA			2 (4%)
FOLLICULAR-CELL CARCINOMA		1 (2%)	1 (2%)
C-CELL ADENOMA	1 (5%)	5 (10%)	4 (8%)
C-CELL CARCINOMA		1 (2%)	
# PANCREATIC ISLETS	(18)	(44)	(48)
ISLET-CELL ADENOMA	1 (6%)	4 (9%)	2 (4%)
ISLET-CELL CARCINOMA			1 (2%)
REPRODUCTIVE SYSTEM			
* PREPUTIAL GLAND	(20)	(50)	(50)
SQUAMOUS CELL CARCINOMA			1 (2%)
# TESTIS	(20)	(49)	(50)
INTERSTITIAL-CELL TUMOR	19 (95%)	48 (98%)	46 (92%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*PLEURA	(20)	(50)	(50)
ALVEOLAR/BRONCHIOLAR CA, INVASIV	1 (5%)		
*TUNICA VAGINALIS	(20)	(50)	(50)
MESOTHELIOMA, NOS		2 (4%)	2 (4%)
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(20)	(50)	(50)
MESOTHELIOMA, NOS		1 (2%)	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	4	11	8
MORIBUND SACRIFICE	1	3	5
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	15	36	37
ANIMAL MISSING			
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	20	50	49
TOTAL PRIMARY TUMORS	42	94	92
TOTAL ANIMALS WITH BENIGN TUMORS	19	49	47
TOTAL BENIGN TUMORS	30	72	69
TOTAL ANIMALS WITH MALIGNANT TUMORS	10	17	19
TOTAL MALIGNANT TUMORS	10	19	21
TOTAL ANIMALS WITH SECONDARY TUMORS#	1		
TOTAL SECONDARY TUMORS	2		
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	2	3	2
TOTAL UNCERTAIN TUMORS	2	3	2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50

INTEGUMENTARY SYSTEM			
* SUBCUT TISSUE	(20)	(50)	(50)
ADENOCARCINOMA, NOS		1 (2%)	
FIBROMA			1 (2%)
NEURILEMOMA			1 (2%)

RESPIRATORY SYSTEM			
# LUNG	(20)	(50)	(50)
SQUAMOUS CELL CARCINOMA			1 (2%)
ADENOCARCINOMA, NOS, METASTATIC		1 (2%)	

HEMATOPOIETIC SYSTEM			
# BRAIN	(20)	(50)	(50)
MALIGNANT RETICULOSIS			1 (2%)
* MULTIPLE ORGANS	(20)	(50)	(50)
MALIGNANT LYMPHOMA, NOS		1 (2%)	6 (12%)
MALIG. LYMPHOMA, UNDIFFER-TYPE	1 (5%)	5 (10%)	9 (18%)
MALIGNANT LYMPHOMA, MIXED TYPE		1 (2%)	
# SPLEEN	(18)	(50)	(50)
HEMANGIOSARCOMA			1 (2%)

CIRCULATORY SYSTEM			
# HEART	(20)	(50)	(50)
ADENOCARCINOMA, NOS, METASTATIC		1 (2%)	

DIGESTIVE SYSTEM			
# CECUM	(19)	(50)	(48)
HEMANGIOSARCOMA		1 (2%)	

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
*PITUITARY	(19)	(48)	(49)
CARCINOMA, NOS	1 (5%)	1 (2%)	
CHROMOPHOBE ADENOMA	8 (42%)	11 (23%)	14 (29%)
*ADRENAL	(20)	(50)	(50)
CORTICAL CARCINOMA	1 (5%)		2 (4%)
PHEOCHROMOCYTOMA		1 (2%)	
*THYROID	(20)	(50)	(49)
FOLLICULAR-CELL CARCINOMA			1 (2%)
C-CELL ADENOMA	4 (20%)	4 (8%)	2 (4%)
C-CELL CARCINOMA		2 (4%)	1 (2%)
*PANCREATIC ISLETS	(20)	(49)	(46)
ISLET-CELL ADENOMA	1 (5%)		
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(20)	(50)	(50)
CARCINOMA, NOS		1 (2%)	
FIBROMA		1 (2%)	1 (2%)
FIBROADENOMA	3 (15%)	8 (16%)	2 (4%)
*CLITORAL GLAND	(20)	(50)	(50)
SQUAMOUS CELL CARCINOMA			1 (2%)
*UTERUS	(20)	(49)	(47)
ENDOMETRIAL STROMAL POLYP	4 (20%)	7 (14%)	3 (6%)
NERVOUS SYSTEM			
*BRAIN	(20)	(50)	(50)
EPENDYMOA		1 (2%)	
SPECIAL SENSE ORGANS			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MUSCULOSKELETAL SYSTEM			
*SKELETAL MUSCLE SARCOMA, NOS	(20)	(50)	(50) 1 (2%)
*MUSCLE OF NECK RHABDOMYOSARCOMA	(20)	(50) 1 (2%)	(50)
BODY CAVITIES			
*MEDIASTINUM SQUAMOUS CELL CARCINOMA, INVASIV	(20)	(50)	(50) 1 (2%)
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH		5	8
MORIBUND SACRIFICE		4	2
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	20	41	40
ANIMAL MISSING			
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	15	33	33
TOTAL PRIMARY TUMORS	23	47	48
TOTAL ANIMALS WITH BENIGN TUMORS	14	21	21
TOTAL BENIGN TUMORS	20	32	24
TOTAL ANIMALS WITH MALIGNANT TUMORS	3	14	21
TOTAL MALIGNANT TUMORS	3	15	24
TOTAL ANIMALS WITH SECONDARY TUMORS#		1	1
TOTAL SECONDARY TUMORS		2	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX B

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MICE ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF THE HISTORY OF ARTS
AND ARCHITECTURE
AND
THE MUSEUM OF ART AND ARCHITECTURE
CHICAGO, ILLINOIS
OFFICE OF THE DIRECTOR
1100 EAST 58TH STREET
CHICAGO, ILLINOIS 60637
TELEPHONE (312) 937-1234
FAX (312) 937-1235
WWW.MUSEUMOFARTANDARCHITECTURE.ORG
MUSEUM OF ART AND ARCHITECTURE
CHICAGO, ILLINOIS
OFFICE OF THE DIRECTOR
1100 EAST 58TH STREET
CHICAGO, ILLINOIS 60637
TELEPHONE (312) 937-1234
FAX (312) 937-1235
WWW.MUSEUMOFARTANDARCHITECTURE.ORG

TABLE B1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		1	
ANIMALS NECROPSIED	20	49	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	49	50
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(49)	(50)
LIPOMA	2 (10%)	3 (6%)	2 (4%)
HEMANGIOSARCOMA		1 (2%)	
NEUROFIBROSARCOMA	1 (5%)		
RESPIRATORY SYSTEM			
#LUNG	(20)	(50)	(48)
HEPATOCELLULAR CARCINOMA, METAST	2 (10%)	2 (4%)	2 (4%)
ALVEOLAR/BRONCHIOLAR ADENOMA	2 (10%)	3 (6%)	6 (13%)
ALVEOLAR/BRONCHIOLAR CARCINOMA	3 (15%)	3 (6%)	2 (4%)
HEMANGIOSARCOMA, METASTATIC			1 (2%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(49)	(50)
MALIGNANT LYMPHOMA, NOS			1 (2%)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE		3 (6%)	3 (6%)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (5%)	1 (2%)	1 (2%)
MAST-CELL SARCOMA		1 (2%)	
*SUBCUT TISSUE	(20)	(49)	(50)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (5%)	1 (2%)	
#SPLEEN	(20)	(49)	(49)
HEMANGIOSARCOMA		1 (2%)	
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE		1 (2%)	
#MESENTERIC L. NODE	(20)	(49)	(49)
HEPATOCELLULAR CARCINOMA, METAST	1 (5%)		
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE			1 (2%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
*SMALL INTESTINE	(20)	(50)	(50)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE	3 (15%)	4 (8%)	
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		1 (2%)	
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER	(20)	(50)	(50)
HEPATOCELLULAR CARCINOMA	10 (50%)	17 (34%)	20 (40%)
HEMANGIOSARCOMA	1 (5%)	1 (2%)	1 (2%)
*ESOPHAGUS	(20)	(50)	(50)
SQUAMOUS CELL CARCINOMA	1 (5%)		
URINARY SYSTEM			
*KIDNEY	(20)	(50)	(50)
HEPATOCELLULAR CARCINOMA, METAST	1 (5%)		
ENDOCRINE SYSTEM			
*THYROID	(20)	(49)	(50)
ADENOCARCINOMA, NOS			2 (4%)
FOLLICULAR-CELL ADENOMA	1 (5%)	3 (6%)	
REPRODUCTIVE SYSTEM			
NONE			
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE/LACRIMAL GLAND	(20)	(49)	(50)
ADENOMA, NOS			4 (8%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	3	7	9
MORIBUND SACRIFICE			
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	17	42	41
ANIMAL MISSING		1	

@ INCLUDES AUTOLYZED ANIMALS

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	16	32	31
TOTAL PRIMARY TUMORS	26	44	43
TOTAL ANIMALS WITH BENIGN TUMORS	4	9	10
TOTAL BENIGN TUMORS	5	9	12
TOTAL ANIMALS WITH MALIGNANT TUMORS	15	27	27
TOTAL MALIGNANT TUMORS	21	35	31
TOTAL ANIMALS WITH SECONDARY TUMORS#	2	2	3
TOTAL SECONDARY TUMORS	4	2	3
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE B2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		3	
ANIMALS NECROPSIED	20	47	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	47	50
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(47)	(50)
FIBROSARCOMA	1 (5%)		1 (2%)
LIPOMA	1 (5%)	1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(47)	(48)
HEPATOCELLULAR CARCINOMA, METAST			1 (2%)
ALVEOLAR/BRONCHIOLAR ADENOMA	2 (10%)	4 (9%)	2 (4%)
ALVEOLAR/BRONCHIOLAR CARCINOMA		2 (4%)	
FIBROSARCOMA, METASTATIC			2 (4%)
OSTEOSARCOMA, METASTATIC			1 (2%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(47)	(50)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE	3 (15%)	2 (4%)	5 (10%)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		6 (13%)	2 (4%)
*SUBCUT TISSUE	(20)	(47)	(50)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE	1 (5%)	2 (4%)	
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	2 (10%)		
*MESENTERIC L. NODE	(20)	(46)	(44)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE		1 (2%)	
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		1 (2%)	
*SMALL INTESTINE	(19)	(46)	(48)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE		1 (2%)	1 (2%)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (5%)		
*THYMUS	(18)	(44)	(42)
FIBROSARCOMA			1 (2%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
MALIG. LYMPHOMA, LYMPHOCYTIC TYPE	1 (6%)	1 (2%)	
MALIG. LYMPHOMA, HISTIOCYTIC TYPE			1 (2%)
CIRCULATORY SYSTEM			
*HEART	(19)	(47)	(48)
FIBROSARCOMA			1 (2%)
DIGESTIVE SYSTEM			
*LIVER	(20)	(47)	(48)
HEPATOCELLULAR CARCINOMA	1 (5%)	2 (4%)	5 (10%)
HEMANGIOSARCOMA	1 (5%)		1 (2%)
*PANCREAS	(17)	(46)	(46)
ADENOCARCINOMA, NOS			1 (2%)
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
*PITUITARY	(19)	(45)	(45)
ADENOMA, NOS		1 (2%)	
*ADRENAL	(20)	(45)	(44)
CORTICAL ADENOMA		1 (2%)	
CORTICAL CARCINOMA			1 (2%)
*THYROID	(20)	(44)	(47)
FOLLICULAR-CELL ADENOMA	1 (5%)		2 (4%)
*PANCREATIC ISLETS	(17)	(46)	(46)
ISLET-CELL ADENOMA			1 (2%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(20)	(47)	(50)
ADENOCARCINOMA, NOS	1 (5%)	1 (2%)	

* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#OVAFY THECOMA	(19)	(47) 1 (2%)	(39)
NERVOUS SYSTEM			
NONF			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
*BONE OSTEOSARCOMA	(20)	(47)	(50) 1 (2%)
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS SARCOMA, NOS HEMANGIOSARCOMA	(20)	(47) 2 (4%)	(50) 1 (2%)
LOWER LEG OSTEOSARCOMA		1	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	5	8	15
MORIBUND SACRIFICE		4	1
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	15	35	34
ANIMAL MISSING		3	
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	15	22	20
TOTAL PRIMARY TUMORS	16	30	27
TOTAL ANIMALS WITH BENIGN TUMORS	4	7	5
TOTAL BENIGN TUMORS	4	8	5
TOTAL ANIMALS WITH MALIGNANT TUMORS	12	20	17
TOTAL MALIGNANT TUMORS	12	22	22
TOTAL ANIMALS WITH SECONDARY TUMORS#			3
TOTAL SECONDARY TUMORS			4
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX C

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN RATS ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

Handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to blurriness and fading.

TABLE C1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
DERMAL INCLUSION CYST	1 (5%)		
INFLAMMATION, SUPPURATIVE	1 (5%)		
GRANULOMA, NOS		1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(48)	(50)
ALVEOLAR MACROPHAGES		1 (2%)	1 (2%)
HYPERPLASIA, ALVEOLAR EPITHELIUM	1 (5%)	5 (10%)	5 (10%)
HEMATOPOIETIC SYSTEM			
#SPLEEN	(19)	(49)	(49)
ECTOPIA	1 (5%)		
INFARCT, FOCAL		1 (2%)	
#PANCREATIC L.NODE	(20)	(50)	(50)
DILATATION, NOS	1 (5%)		
#MESENTERIC L. NODE	(20)	(50)	(50)
DILATATION, NOS		12 (24%)	32 (64%)
CIRCULATORY SYSTEM			
#HEART/ATRIUM	(20)	(50)	(50)
THROMBOSIS, NOS		1 (2%)	1 (2%)
#MYOCARDIUM	(20)	(50)	(50)
FIBROSIS	8 (40%)	2 (4%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
FIBROSIS, FOCAL			1 (2%)
DIGESTIVE SYSTEM			
*LIVER	(20)	(50)	(49)
NECROSIS, FOCAL		1 (2%)	
METAMORPHOSIS FATTY		2 (4%)	1 (2%)
HEPATOCYTOMEGALY	5 (25%)	10 (20%)	10 (20%)
HYPERPLASIA, FOCAL	1 (5%)	1 (2%)	
*BILE DUCT	(20)	(50)	(50)
HYPERPLASIA, NOS	14 (70%)	2 (4%)	
HYPERPLASIA, FOCAL		1 (2%)	
*PANCREAS	(18)	(44)	(48)
FIBROSIS, FOCAL	1 (6%)		
PERIARTERITIS	3 (17%)	1 (2%)	1 (2%)
*PANCREATIC ACINUS	(18)	(44)	(48)
ATROPHY, NOS	3 (17%)	3 (7%)	4 (8%)
*STOMACH	(19)	(50)	(50)
ULCER, NOS			1 (2%)
ULCER, FOCAL		1 (2%)	2 (4%)
NECROSIS, FOCAL			1 (2%)
*LARGE INTESTINE	(20)	(49)	(48)
NEMATODIASIS		1 (2%)	
HYPERPLASIA, LYMPHOID	1 (5%)		
URINARY SYSTEM			
*KIDNEY	(20)	(50)	(50)
CYST, NOS			1 (2%)
INFLAMMATION, CHRONIC	18 (90%)	41 (82%)	45 (90%)
PERIARTERITIS			1 (2%)
HYPERPLASIA, TUBULAR CELL			1 (2%)
*KIDNEY/CORTEX	(20)	(50)	(50)
NECROSIS, FOCAL		1 (2%)	
*KIDNEY/PELVIS	(20)	(50)	(50)
DILATATION, NOS			1 (2%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#URINARY BLADDER HYPERPLASIA, EPITHELIAL	(18)	(46)	(50) 2 (4%)
ENDOCRINE SYSTEM			
#PITUITARY HEMORRHAGE	(19) 1 (5%)	(49) 3 (6%)	(48) 2 (4%)
#ADRENAL METAPLASIA, OSSEOUS	(20)	(50) 1 (2%)	(50)
#ADRENAL CORTEX HYPERPLASIA, FOCAL	(20)	(50)	(50) 1 (2%)
#ADRENAL MEDULLA HYPERPLASIA, FOCAL	(20)	(50) 3 (6%)	(50) 5 (10%)
#THYROID HYPERPLASIA, C-CELL	(20) 8 (40%)	(49) 10 (20%)	(50) 12 (24%)
#PARATHYROID HYPERPLASIA, NOS	(18)	(46)	(42) 1 (2%)
#PANCREATIC ISLETS HYPERPLASIA, NOS	(18) 1 (6%)	(44)	(48)
REPRODUCTIVE SYSTEM			
*PREPUTIAL GLAND CYST, NOS	(20)	(50)	(50) 1 (2%)
INFLAMMATION, GRANULOMATOUS HYPERPLASIA, NOS		1 (2%) 2 (4%)	
#PROSTATE INFLAMMATION, SUPPURATIVE HYPERPLASIA, EPITHELIAL	(18) 2 (11%)	(47) 8 (17%)	(49) 17 (35%) 1 (2%)
#TESTIS PERIARTERITIS	(20)	(49)	(50) 1 (2%)
*EPIDIDYMIS GRANULOMA, SPERMATIC	(20)	(50) 1 (2%)	(50)

* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
NERVOUS SYSTEM			
# BRAIN	(20)	(49)	(49)
HEMORRHAGE	2 (10%)	3 (6%)	
CORPORA AMYLACEA			1 (2%)
SPECIAL SENSE ORGANS			
* EYE	(20)	(50)	(50)
CATARACT	3 (15%)		
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
* MESENTERY	(20)	(50)	(50)
PERIARTERITIS		1 (2%)	
ALL OTHER SYSTEMS			
ADIPOSE TISSUE			
INFLAMMATION, FOCAL		3	2
SPECIAL MORPHOLOGY SUMMARY			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
#LUNG	(20)	(50)	(50)
ALVEOLAR MACROPHAGES	1 (5%)	6 (12%)	3 (6%)
HYPERPLASIA, ALVEOLAR EPITHELIUM		3 (6%)	6 (12%)
HEMATOPOIETIC SYSTEM			
#BONE MARROW	(20)	(49)	(49)
HYPERPLASIA, HEMATOPOIETIC			1 (2%)
#SPLEEN	(18)	(50)	(50)
NECROSIS, FOCAL		1 (2%)	
HEMATOPOIESIS		3 (6%)	1 (2%)
#MESENTERIC L. NODE	(20)	(50)	(50)
DILATATION, NOS		5 (10%)	18 (36%)
INFLAMMATION, PYOGRANULOMATOUS		1 (2%)	
NECROSIS, NOS		1 (2%)	
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#LIVER	(20)	(50)	(49)
NECROSIS, FOCAL		2 (4%)	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
METAMORPHOSIS FATTY	2 (10%)	2 (4%)	
HEPATOCYTOMEGALY	3 (15%)	5 (10%)	10 (20%)
HYPERPLASIA, FOCAL	13 (65%)	28 (56%)	12 (24%)
HEMATOPOIESIS		1 (2%)	
*BILE DUCT	(20)	(50)	(50)
HYPERPLASIA, NOS		1 (2%)	1 (2%)
#PANCREATIC ACINUS	(20)	(49)	(46)
ATROPHY, NOS	1 (5%)	5 (10%)	3 (7%)
#LARGE INTESTINE	(19)	(50)	(48)
NEMATODIASIS		1 (2%)	1 (2%)
URINARY SYSTEM			
*KIDNEY	(20)	(50)	(50)
PYELONEPHRITIS SUPPURATIVE		1 (2%)	
INFLAMMATION, CHRONIC	8 (40%)	25 (50%)	26 (52%)
PERIARTERITIS	1 (5%)		
*KIDNEY/PELVIS	(20)	(50)	(50)
HYPERPLASIA, EPITHELIAL			1 (2%)
#URINARY BLADDER	(19)	(48)	(46)
HYPERPLASIA, EPITHELIAL		1 (2%)	
ENDOCRINE SYSTEM			
*PITUITARY	(19)	(48)	(49)
CYST, NOS		1 (2%)	
*ADRENAL CORTEX	(20)	(50)	(50)
NECROSIS, NOS		1 (2%)	
NECROSIS, FOCAL		1 (2%)	
CYTOMEGALY		1 (2%)	
*ADRENAL MEDULLA	(20)	(50)	(50)
EOSINOPHILIC INFILTRATE		1 (2%)	
HYPERPLASIA, FOCAL		1 (2%)	
*THYROID	(20)	(50)	(49)
CYSTIC FOLLICLES			1 (2%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
HYPERPLASIA, C-CELL	11 (55%)	22 (44%)	22 (45%)
HYPERPLASIA, FOLLICULAR-CELL			2 (4%)
#PARATHYROID	(18)	(45)	(43)
HYPERPLASIA, NOS			1 (2%)
REPRODUCTIVE SYSTEM			
*MAMMARY LOBULE	(20)	(50)	(50)
HYPERPLASIA, NOS	1 (5%)	1 (2%)	
HYPERPLASIA, FOCAL			1 (2%)
#UTERUS/ENDOMETRIUM	(20)	(49)	(47)
HYPERPLASIA, NOS			1 (2%)
HYPERPLASIA, CYSTIC	1 (5%)	1 (2%)	1 (2%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE	(20)	(50)	(50)
CATARACT		1 (2%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(20)	(50)	(50)
BACTERIAL SEPTICEMIA		1 (2%)	
ADIPOSE TISSUE			
INFLAMMATION, FOCAL		2	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LDW DDSE	HIGH DDSE
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED			1
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX D

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MICE ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

TABLE D1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		1	
ANIMALS NECROPSIED	20	49	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	49	50
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(49)	(50)
ALOPECIA		2 (4%)	2 (4%)
RESPIRATORY SYSTEM			
NONE			
HEMATOPOIETIC SYSTEM			
*SPLEEN	(20)	(49)	(49)
HYPERPLASIA, LYMPHOID	2 (10%)	1 (2%)	3 (6%)
HEMATOPOIESIS	6 (30%)	5 (10%)	1 (2%)
*MANDIBULAR L. NODE	(20)	(49)	(49)
HYPERPLASIA, LYMPHOID			1 (2%)
*MESENTERIC L. NODE	(20)	(49)	(49)
HYPERPLASIA, RETICULUM CELL		1 (2%)	1 (2%)
HYPERPLASIA, LYMPHOID			1 (2%)
HEMATOPOIESIS	4 (20%)		1 (2%)
*THYMUS	(20)	(47)	(48)
ATROPHY, NOS			2 (4%)
HYPERPLASIA, LYMPHOID	1 (5%)	1 (2%)	3 (6%)
CIRCULATORY SYSTEM			
*MYOCARDIUM	(20)	(50)	(50)
INFLAMMATION, NOS		1 (2%)	2 (4%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
DIGESTIVE SYSTEM			
#LIVER	(20)	(50)	(50)
NECROSIS, NOS		1 (2%)	
NECROSIS, FOCAL	1 (5%)		2 (4%)
HYPERPLASIA, NODULAR	1 (5%)	3 (6%)	5 (10%)
ANGIECTASIS	1 (5%)		
HEMATOPOIESIS	1 (5%)		
*GALLBLADDER	(20)	(49)	(50)
CALCULUS, NOS	1 (5%)		
*BILE DUCT	(20)	(49)	(50)
INFLAMMATION, NOS	2 (10%)		3 (6%)
#PANCREAS	(20)	(47)	(49)
CYSTIC DUCTS			1 (2%)
ATROPHY, NOS		1 (2%)	
#STOMACH	(20)	(49)	(50)
INFLAMMATION, NOS	2 (10%)		
#PEYERS PATCH	(20)	(50)	(50)
HYPERPLASIA, NOS			1 (2%)
URINARY SYSTEM			
#KIDNEY	(20)	(50)	(50)
HYDRONEPHROSIS			1 (2%)
INFLAMMATION, INTERSTITIAL	1 (5%)	2 (4%)	
INFLAMMATION, CHRONIC			1 (2%)
PERIVASCULITIS			1 (2%)
INFARCT, NOS		1 (2%)	
ENDOCRINE SYSTEM			
#THYROID	(20)	(49)	(50)
CYSTIC FOLLICLES	1 (5%)		
FOLLICULAR CYST, NOS		1 (2%)	2 (4%)
#PARATHYROID	(15)	(33)	(34)
CYST, NOS			1 (3%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
*PANCREATIC ISLETS HYPERPLASIA, NOS	(20) 3 (15%)	(47) 3 (6%)	(49) 2 (4%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND HYPERPLASIA, CYSTIC	(20) 1 (5%)	(49)	(50)
*PREPUTIAL GLAND CYST, NOS	(20)	(49) 2 (4%)	(50) 2 (4%)
*SEMINAL VESICLE CAST, NOS	(20) 6 (30%)	(49) 11 (22%)	(50) 10 (20%)
*TESTIS ATROPHY, NOS	(20)	(48)	(50) 1 (2%)
NERVOUS SYSTEM			
*BRAIN MINERALIZATION	(20)	(50) 4 (8%)	(49) 8 (16%)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*PLEURA INFLAMMATION, NOS	(20) 1 (5%)	(49)	(50)
ALL OTHER SYSTEMS			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	1	10	7
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE
ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		3	
ANIMALS NECROPSIED	20	47	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	47	50
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(47)	(50)
ALOPECIA		3 (6%)	3 (6%)
RESPIRATORY SYSTEM			
*LUNG	(20)	(47)	(48)
ATELECTASIS	1 (5%)		2 (4%)
CONGESTION, NOS		3 (6%)	
HEMATOPOIETIC SYSTEM			
*SPLEEN	(19)	(47)	(46)
HYPERPLASIA, LYMPHOID		4 (9%)	4 (9%)
HEMATOPOIESIS	8 (42%)	8 (17%)	3 (7%)
*LYMPH NODE	(20)	(46)	(44)
CYST, NOS			1 (2%)
*MESENTERIC L. NODE	(20)	(46)	(44)
CYST, NOS	1 (5%)	1 (2%)	
HYPERPLASIA, LYMPHOID	1 (5%)		3 (7%)
*THYMUS	(18)	(44)	(42)
HYPERPLASIA, LYMPHOID	2 (11%)		1 (2%)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER	(20)	(47)	(48)
NECROSIS, FOCAL		2 (4%)	2 (4%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
HYPERPLASIA, NODULAR HEMATOPOIESIS	3 (15%) 1 (5%)	7 (15%)	4 (8%)
*HEPATIC CAPSULE INFLAMMATION, NOS	(20) 1 (5%)	(47)	(48)
*GALLBLADDER HYPERPLASIA, NOS	(20) 1 (5%)	(47)	(50)
*BILE DUCT INFLAMMATION, NOS	(20) 2 (10%)	(47) 5 (11%)	(50) 1 (2%)
*PANCREAS CYSTIC DUCTS ATPOPHY, NOS	(17) 1 (6%)	(46) 1 (2%)	(46)
*STOMACH INFLAMMATION, NOS	(20) 3 (15%)	(47) 2 (4%)	(48) 1 (2%)
*SMALL INTESTINE POLYP	(19) 1 (5%)	(46)	(48)
*PEYERS PATCH HYPERPLASIA, NOS	(19)	(46)	(48) 1 (2%)
URINARY SYSTEM			
*KIDNEY PERIVASCULITIS HYPERPLASIA, LYMPHOID	(20) 1 (5%)	(47) 1 (2%)	(48)
ENDOCRINE SYSTEM			
*PITUITARY ANGIECTASIS	(19) 1 (5%)	(45)	(45)
*THYROID CYSTIC FOLLICLES	(20)	(44) 1 (2%)	(47) 5 (11%)
*PANCREATIC ISLETS HYPERPLASIA, NOS	(17)	(46) 1 (2%)	(46)
REPRODUCTIVE SYSTEM			
*UTERUS INFLAMMATION, NOS	(20)	(47)	(44) 1 (2%)

* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
POLYPOID HYPERPLASIA	2 (10%)		
#UTERUS/ENDOMETRIUM	(20)	(47)	(44)
CYST, NOS	9 (45%)	21 (45%)	22 (50%)
HYPERPLASIA, NOS			1 (2%)
#OVARY	(19)	(47)	(39)
CYST, NOS	5 (26%)	11 (23%)	5 (13%)
NERVOUS SYSTEM			
#BRAIN	(20)	(47)	(48)
MINIFALIZATION	2 (10%)	9 (19%)	6 (13%)
INFLAMMATION, FOCAL		1 (2%)	
SPECIAL SENSE ORGANS			
*EYE/LACRIMAL GLAND	(20)	(47)	(50)
CYST, NOS		1 (2%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	1	6	8
ANIMAL MISSING/NO NECROPSY		1	
AUTO/NECROPSY/HISTO PERF			1
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX E

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS
IN RATS ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

THE JOURNAL OF THE
ROYAL ANTHROPOLOGICAL INSTITUTE

Table El. Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Integumentary System: Fibroma of the Subcutaneous Tissue (b)	0/20 (0)	2/50 (4)	3/50 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		Infinite	Infinite
Upper Limit		0.123	0.250
		Infinite	Infinite
Weeks to First Observed Tumor	--	104	107
<hr/>			
Hematopoietic System: Lymphoma (b)	9/20 (45)	15/50 (30)	13/50 (26)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.667	0.578
Upper Limit		0.346	0.289
		1.478	1.316
Weeks to First Observed Tumor	91	81	77

Table E1. Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Liver: Neoplastic Nodule (b)	2/20 (10)	0/50 (0)	0/49 (0)
P Values (c,d)	P = 0.025(N)	N.S.	N.S.
Departure from Linear Trend (e)	P = 0.043		
Relative Risk (f)			
Lower Limit		0.000	0.000
Upper Limit		0.000	0.000
		1.345	1.372
Weeks to First Observed Tumor	98	--	--
Pituitary: Adenoma or Carcinoma, NOS (b)	4/19 (21)	0/49 (0)	1/48 (2)
P Values (c,d)	P = 0.009(N)	P = 0.005(N)	P = 0.020(N)
Departure from Linear Trend (e)	P = 0.004		
Relative Risk (f)			
Lower Limit		0.000	0.099
Upper Limit		0.000	0.002
		0.413	0.932
Weeks to First Observed Tumor	102	--	103

Table E1. Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Pituitary: Chromophobe Adenoma (b)	1/19 (5)	7/49 (14)	6/48 (13)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		2.714	2.375
Upper Limit		0.393	0.324
		119.544	106.788
Weeks to First Observed Tumor	107	104	107
Adrenal: Pheochromocytoma (b)	2/20 (10)	4/50 (8)	4/50 (8)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.800	0.800
Upper Limit		0.128	0.128
		8.436	8.436
Weeks to First Observed Tumor	107	57	104

Table E1. Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

(continued)				
<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>	
Pituitary: Chromophobe Adenoma, Adenoma, NOS or Carcinoma (b)	5/19 (26)	7/49 (14)	7/48 (15)	
P Values (c,d)	N.S.	N.S.	N.S.	
Relative Risk (f)				
Lower Limit		0.543	0.554	
Upper Limit		0.176	0.180	
		1.959	1.997	
Weeks to First Observed Tumor	102	104	103	
Thyroid: Follicular-cell Adenoma or Carcinoma (b)	0/20 (0)	1/49 (2)	3/50 (6)	
P Values (c,d)	N.S.	N.S.	N.S.	
Relative Risk (f)				
Lower Limit		Infinite	Infinite	
Upper Limit		0.023	0.250	
		Infinite	Infinite	
Weeks to First Observed Tumor	--	107	81	

Table E1 Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

Topography: Morphology	Matched Control	Low Dose	High Dose
Thyroid: C-cell Adenoma or Carcinoma (b)	1/20 (5)	6/49 (12)	4/50 (8)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		2.449	1.600
Upper Limit		0.332	0.175
		110.166	77.169
Weeks to First Observed Tumor	107	104	107
Pancreatic Islets: Islet-cell Adenoma or Carcinoma (b)	1/18 (6)	4/44 (9)	3/48 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.636	1.125
Upper Limit		0.181	0.100
		78.690	57.811
Weeks to First Observed Tumor	107	107	107

Table El. Analyses of the Incidence of Primary Tumors in Male Rats
Administered Piperonyl Butoxide in the Diet (a)

(continued)			
<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Testis: Interstitial-cell Tumor (b)	19/20 (95)	48/49 (98)	46/50 (92)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.031	0.968
Upper Limit		0.961	0.902
		1.109	1.174
Weeks to First Observed Tumor	96	78	77

(a) Dosed groups received 5,000 or 10,000 ppm.

(b) Number of tumor-bearing animals/number of animals examined at site (percent).

(c) Beneath the incidence of tumors in the matched-control group is the probability level for the Cochran-Armitage test when P less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P less than 0.05; otherwise, not significant (N.S.) is indicated.

(d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.

(e) The probability level for departure from linear trend is given when P less than 0.05 for any comparison.

(f) The 95% confidence interval of the relative risk between each dosed group and the control group.

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats Administered Piperonyl Butoxide in the Diet (a)

<u>Topography:</u>	<u>Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
	Hematopoietic System: Lymphoma (b)	1/20 (5)	7/50 (14)	15/50 (30)
P Values (c,d)		P = 0.007	N.S.	P = 0.020
Relative Risk (f)				
Lower Limit			2.800	6.000
Upper Limit			0.403	1.048
			123.407	245.704
Weeks to First Observed Tumor		107	94	73
	Pituitary: Chromophobe Adenoma (b)	8/19 (42)	11/48 (23)	14/49 (29)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			0.544	0.679
Upper Limit			0.251	0.336
			1.348	1.606
Weeks to First Observed Tumor		107	107	105

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats Administered Piperonyl Butoxide in the Diet (a)

(continued)			
<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Thyroid: C-cell Adenoma or Carcinoma (b)	4/20 (20)	6/50 (12)	3/49 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.600	0.306
Upper Limit		0.164	0.050
		2.659	1.675
Weeks to First Observed Tumor	107	107	105
Mammary Gland: Fibroadenoma (b)	3/20 (15)	8/50 (16)	2/50 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.067	0.267
Upper Limit		0.295	0.024
		5.813	2.190
Weeks to First Observed Tumor	107	107	105

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography:</u>	<u>Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Uterus:	Endometrial Stromal Polyp (b)	4/20 (20)	7/49 (14)	3/47 (6)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			0.714	0.319
Upper Limit			0.211	0.052
			3.052	1.743
Weeks to First Observed Tumor		107	107	107

(a) Dosed groups received 5,000 or 10,000 ppm.

(b) Number of tumor-bearing animals/number of animals examined at site (percent).

(c) Beneath the incidence of tumors in the matched-control group is the probability level for the Cochran-Armitage test when P less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P less than 0.05; otherwise, not significant (N.S.) is indicated.

(d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.

(e) The probability level for departure from linear trend is given when P less than 0.05 for any comparison.

(f) The 95% confidence interval of the relative risk between each dosed group and the control group.

APPENDIX F

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS IN MICE ADMINISTERED PIPERONYL BUTOXIDE IN THE DIET

These figures are given as a guide only
and are not intended to be used as a basis for
any other calculations.

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Administered Piperonyl Butoxide in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Integumentary System: Lipoma of the Subcutaneous Tissue (b)	2/20 (10)	3/49 (6)	2/50 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.612	0.400
Upper Limit		0.078	0.032
		6.996	5.277
Weeks to First Observed Tumor	112	112	112
Lung: Alveolar/Bronchiolar Carcinoma (b)	3/20 (15)	3/50 (6)	2/48 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.400	0.278
Upper Limit		0.060	0.025
		2.802	2.278
Weeks to First Observed Tumor	103	105	112

Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice
Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung: Alveolar/Bronchiolar Adenoma or Carcinoma (b)	5/20 (25)	6/50 (12)	8/48 (17)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.480	0.667
Upper Limit		0.143	0.227
		1.807	2.338
Weeks to First Observed Tumor	103	105	112
Hematopoietic System: Lymphoma (b)	4/20 (20)	10/49 (20)	6/50 (12)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.020	0.600
Upper Limit		0.346	0.164
		4.068	2.659
Weeks to First Observed Tumor	112	91	90

Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice
Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
All Sites: Hemangiosarcoma (b)	1/20 (5)	3/49 (6)	1/50 (2)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.224	0.400
Upper Limit		0.108	0.005
		62.958	30.802
Weeks to First Observed Tumor	112	105	112
Liver: Hepatocellular Carcinoma (b)	10/20 (50)	17/50 (34)	20/50 (40)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.680	0.800
Upper Limit		0.379	0.462
		1.403	1.604
Weeks to First Observed Tumor	88	83	99

Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice
Administered Piperonyl Butoxide in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Thyroid: Follicular-cell Adenoma (b)	1/20 (5)	3/49 (6)	0/50 (0)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.224	0.000
Upper Limit		0.108	0.000
		62.958	7.475
Weeks to First Observed Tumor	112	112	--
Lacrimal Gland: Adenoma, NOS (b)	0/20 (0)	0/49 (0)	4/50 (8)
P Values (c,d)	P = 0.022	N.S.	N.S.
Relative Risk (f)			
Lower Limit		--	Infinite
Upper Limit		--	0.386
		--	Infinite
Weeks to First Observed Tumor	--	--	112

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice
Administered Piperonyl Butoxide in the Diet (a)

(continued)

- (a) Dosed groups received time-weighted average doses of 1,036 or 2,804 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the matched-control group is the probability level for the Cochran-Armitage test when P less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.
- (e) The probability level for departure from linear trend is given when P less than 0.05 for any comparison.
- (f) The 95% confidence interval of the relative risk between each dosed group and the control group.

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Administered Piperonyl Butoxide in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung: Alveolar/Bronchiolar Adenoma or Carcinoma (b)	2/20 (10)	6/47 (13)	2/48 (4)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.277	0.417
Upper Limit		0.258	0.033
		12.285	5.490
Weeks to First Observed Tumor	111	98	112
Hematopoietic System: Lymphoma (b)	8/20 (40)	14/47 (30)	9/50 (18)
P Values (c,d)	P = 0.030(N)	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.745	0.450
Upper Limit		0.364	0.190
		1.765	1.174
Weeks to First Observed Tumor	82	45	73

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Administered Piperonyl Butoxide in the Diet (a)

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>
Liver:	Hepatocellular Carcinoma (b)	1/20 (5)	2/47 (4)
P Values (c,d)		N.S.	N.S.
Relative Risk (f)			
Lower Limit			0.851
Upper Limit			0.048
			49.164
Weeks to First Observed Tumor		112	111
			107

- (a) Dosed groups received time-weighted average doses of 1,036 or 2,804 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the matched-control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.
- (e) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.
- (f) The 95% confidence interval of the relative risk between each dosed group and the control group.

Review of the Bioassay of Piperonyl Butoxide* for Carcinogenicity
by the Data Evaluation/Risk Assessment Subgroup of the
Clearinghouse on Environmental Carcinogens

August 31, 1978

The Clearinghouse on Environmental Carcinogens was established in May, 1976, in compliance with DHEW Committee Regulations and the Provisions of the Federal Advisory Committee Act. The purpose of the Clearinghouse is to advise the Director of the National Cancer Institute (NCI) on its bioassay program to identify and to evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, State health officials, and quasi-public health and research organizations. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of reports prepared on NCI-sponsored bioassays of chemicals studied for carcinogenicity. It is in this context that the below critique is given on the bioassay of Piperonyl Butoxide for carcinogenicity.

The primary reviewer said that the staff concluded that Piperonyl Butoxide was not carcinogenic in rats or mice, under the conditions of test. He briefly described the experimental design and noted the increased incidence of lymphomas observed among treated female rats. He said that the neoplasms were not regarded as significant when compared to historical controls. The primary reviewer concluded that the study was adequate and his recommendation was approved unanimously that the report be accepted as written.

Members present were:

Arnold L. Brown (Chairman), University of Wisconsin Medical School
Joseph Highland, Environmental Defense Fund
Michael Shimkin, University of California at San Diego
Louise Strong, University of Texas Health Sciences Center

* Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.

10 Center Drive
Bethesda, MD 20892-1150
301-496-1080

4 0130 7595



DHEW Publication No. (NIH) 79-1375